

**RADIOLABELED ANNEXINS**Cross-Reference to Related Applications

5 This application is a continuation-in-part of U.S. Patent Application No. 08/351,635, filed December 7, 1994, which is in turn a continuation-in-part of pending U.S. Patent Application No. 08/261,064, filed June 16, 1994, which is in turn a continuation-in-part of pending U.S. Patent Application No. 08/185,660, filed January 24, 1994.

Technical Field

10 The present invention is directed to radiolabeled annexins, including annexin-hexose conjugates such as hexose moiety conjugates, and components thereof. The present invention is also generally directed to modifications of the annexin components for the use in the preparation of radiolabeled annexins. Also contemplated by the present invention are imaging protocols which involve the administration of, for example, a radiolabeled annexin-hexose moietyconjugate. The annexin component of the conjugate serves to deliver the radiolabeled active component of the conjugate to vascular thrombi target sites. The hexose component of the conjugate facilitates rapid elimination of the radiolabeled annexin-hexose moiety conjugates from the circulation of a recipient.

15

20

25

### Background of the Invention

When patients present with chest pain, palpitations or any other symptoms of coronary trauma or disease, the presence of vascular thrombi in the heart is a potential significant complicating factor for treatment. If a medical practitioner could non-invasively determine whether one or more vascular thrombi were present and, if present, the location of those vascular thrombi, better evaluation of treatment options would be possible. Furthermore, if a medical practitioner could determine that no vascular thrombi were present, thereby eliminating a potential complication in treatment, cardiac conditions could be treated more safely and effectively.

Most present techniques for determining the presence of vascular thrombi are invasive and/or cumbersome, and/or fail to detect such thrombi with good sensitivity and specificity. Thus, an imaging agent useful for non-invasive vascular thrombi imaging is desirable.

Annexins are a class of proteins that are characterized by calcium-mediated binding to anionic phospholipids. Anionic phospholipids are about 20-fold more highly associated with activated platelets than quiescent platelets, and activated platelets are associated with vascular thrombi.

Radioiodinated annexin V has been shown to localize to vascular thrombi in vivo, but has suboptimal imaging characteristics, possibly due to the pronounced beta

phase of blood clearance owing to possible transiodination and/or metabolic degradation with reincorporation into serum macromolecules or non-target tissues. Free radioactive iodine or iodine-containing metabolic degradation products exposed non-target tissues, especially the thyroid gland, to radioactivity. Iodine radioisotopes I-123, I-124 and I-131 with imagable photons suffer from various drawbacks. Iodine-131 has particulate emission and its gamma emission is too high in energy for optimal imaging. Iodine-124 emits positrons and high energy gamma photons. Finally, I-iodine-123 radiolabel with superior imaging properties is expensive and difficult to obtain and is not therefore practical for wide spread use. Consequently, improved radiolabeled annexin compounds are desirable.

In addition, conventional imaging and therapy are often plagued by the problem that the generally attainable targeting ratio (ratio of administered dose localizing to target versus administered dose circulating in blood, or ratio of administered dose localizing to target versus administered dose migrating to bone marrow) is low. Improvement in targeting ratio is also sought.

Thus, for the foregoing reasons there is still a need for a thrombus imaging product with increased sensitivity to image small thrombi, such as carotid thrombi or intracardiac thrombi and those present in coronary arteries after angioplasty or during myocardial infarctions or in cerebral arteries during stroke. Development of a

radiolabeled thrombus imaging agent capable of detecting intracoronary thrombi would represent a breakthrough product of great clinical and commercial significance. The present invention fulfills this need and provides further related advantages.

#### Summary of the Invention

The present invention provides radiolabeled annexin-hexose moiety conjugates and methods of making and using the same. The present invention further provides an annexin component that is altered in such a way as to provide an accessible sulfhydryl group and/or other amino acid groups for derivatization so as to serve to improve the imaging properties of annexin. Annexin-containing conjugates of the present invention are suitable for radiolabeling with a diagnostic imaging agent. Preferred conjugates of the present invention include:

- an annexin;
- a hexose moiety; and
- an  $N_2S_2$  chelating compound associated with the annexin.

Also provided by the present invention are radiolabeled annexin-hexose moiety-conjugates which include:

- an annexin;
- a hexose moiety;
- an  $N_2S_2$  chelating compound associated with the annexin; and

a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention includes:

5 a modified annexin, wherein the modification provides an accessible sulfhydryl group; and

a hexose moiety recognized by a mammalian liver receptor, wherein the hexose moiety is conjugated to the annexin.

10 Another preferred conjugate of the present invention includes:

a modified annexin, wherein the modification provides an accessible sulfhydryl group;

15 a hexose moiety recognized by a mammalian liver receptor; and

20 a  $N_xS_y$  chelating compound, wherein the hexose moiety is conjugated to the modified annexin directly or via the chelating compound and the chelating compound is conjugated to the modified annexin directly or via the hexose moiety.

Another preferred conjugate of the present invention includes:

an annexin multimer;

25 a hexose moiety recognized by a mammalian liver receptor; and

a  $N_xS_y$  chelating compound, wherein the hexose moiety is conjugated to the multimer directly or via the chelating compound and the chelating compound is

conjugated to the multimer directly or via the hexose moiety.

Another preferred conjugate of the present invention includes:

5           an annexin; and  
          an esterase-sensitive  $N_xS_y$  chelating compound  
conjugated to the annexin.

Another preferred conjugate of the present invention includes:

10           a modified annexin, wherein the modification provides  
accessible sulfhydryl group; and

          a  $N_xS_y$  chelating compound conjugated to the annexin.

Another preferred conjugate of the present invention includes:

15           a modified annexin, wherein the modification provides  
an accessible sulfhydryl group; and

          an esterase-sensitive  $N_xS_y$  chelating compound  
conjugated to the annexin.

Another preferred conjugate of the present invention includes:

20           an annexin multimer; and

          a  $N_xS_y$  chelating compound conjugated to the annexin.

Another preferred conjugate of the present invention includes:

25           an annexin multimer; and

          an esterase-sensitive  $N_xS_y$  chelating compound  
conjugated to the annexin.

It will be understood by one of ordinary skill in the

art that where a chelating compound is a component of the conjugate, any one of the above preferred conjugates further includes a radionuclide complexed by the chelating compound.

5 A preferred annexin for use in the present invention is annexin V. A modified annexin is extended on the C-terminus or N-terminus by a number of amino acids. A preferred modified annexin for use in the present invention is extended on the N-terminus. This addition of the extension results in a modification to the annexin whereby an accessible sulfhydryl group is provided for. Thus, the modified annexin can be conjugated to a hexose moiety, such as a hexose cluster, (e.g., N-acetylgalactos-amine cluster; or provide for an endogenous radiolabel chelation site, thus eliminating the need for intermediate production of an amine-directed active ester containing the radiolabel; or provide for multimer production.

10 The modified annexin component of the conjugate allows for rapid binding of the radiolabel to the annexin, thus providing for a simplified labeling procedure for clinical use. In some embodiments of the present invention, the modification to the annexin provides for acceleration of blood clearance thereby reducing the background radioactivity of the blood pool.

20 Also, in some other embodiments of the present invention, an esterase-sensitive chelating compound is utilized to provide for accelerated excretion of

radioactivity from the liver, thereby reducing background radioactivity adjacent to the heart and lungs.

Preferred hexose moieties for use in the present invention include hexose clusters and incorporate a multiple of 4 galactoses. Other preferred hexose clusters for use in the present invention incorporate a multiple of 3 galactoses.

The hexose moiety component may bound to the annexin component and the chelating compound component via a trifunctional linker, such as lysine. An extender may be employed between the hexose moiety and the trifunctional linker to promote bioavailability of the hexose moiety. This embodiment of the present invention is favored if the chelating compound is characterized by a single functional group available and suitable for conjugation.

If the chelating compound component is characterized by greater than one functional group available and suitable for conjugation to other conjugate components, a structure such as the following is preferably employed:  
hexose moiety - bifunctional linker - chelating compound - bifunctional linker - annexin.

Preferred diagnostic radionuclides for use in the practice of the present invention are Tc-99m, Re-186 and Re-188, with Tc-99m being especially preferred. Vascular thrombi located in or near the heart are especially amenable to imaging in accordance with the present invention.

These and other aspects of the present invention will

become evident upon reference to the following description and attached drawings.

#### Brief Description of the Drawings

5           Fig. 1 schematically represents a method of radiolabeling annexin V.

Fig. 2 shows the blood clearance of Tc-99m-annexin V (o) and of Tc-99m-annexin V-galactose ( $\Delta$ )

Fig. 3 schematically shows the synthesis of N,N'-bis(2-disulfidyl-4-carbonylphenyl)-1,3-propyldiamine.

Fig. 4 schematically shows the synthesis of N,N'-bis(disulfidyl-4-methylphenyl)-gamma,gamma'-diamino-isovalerate N-hydroxysuccinimide.

Figs. 5a and 5b schematically show the synthesis of an eight galactose-containing galactose cluster.

Fig. 6 schematically shows the synthesis of an extended eight galactose-containing galactose cluster.

Fig. 7 schematically represents a pET-12a plasmid Map.

20           Fig. 8 schematically represents pET-12a-PAP1,3/7/94, clone 1.

#### Detailed Description of the Invention

25           Prior to setting forth the invention, it may be helpful to set forth definitions of certain terms to be used within the disclosure.

Annexin: A class of compounds characterized by the

ability to bind with high affinity to membrane lipids in the presence of millimolar concentrations of calcium. Annexins have been shown to exhibit anti-coagulatory effects that are mediated by the binding of annexins to negatively charged surface phospholipids (e.g., on activated platelets). This annexin-phospholipid binding is believed to block the activation of clotting factors by such negatively charged surface phospholipids. Prior to the recognition of the annexin class of molecules, members thereof were also referred to in the literature as placental anticoagulant proteins (e.g., PAP-1, 2, 3 and 4), lipocortins, calpactins, vascular anti-coagulant (alpha and beta), calphobindin I, placental protein 4 (PP4), endonexin II, anchorin CII, calcium-dependent phospholipid binding protein, and the like. See Crumpton et al., Nature 345:212, 1990. Annexin-V is a prototypical annexin molecule used in the description of the present invention. The term annexin includes native annexin purified from natural sources such as e.g. human placenta, or annexin molecules containing a native sequence produced through e.g. genetic engineering, recombinant, or other means. The term annexin further includes modified annexins as defined below, derived from or produced by any source.

Modified Annexin: An annexin molecule wherein the native sequence or molecule is altered in such a way without materially altering the membrane binding affinity of the annexin. Such annexins can be produced by

chemical, genetic engineering or recombinant techniques as know to those of ordinary skill in the art. The modification can include a modification of the sequence through the addition of several amino acid residues, and/or an addition/deletion of an amino acid at a single site on the native or genetically engineered sequence. For example, the annexin can be modified at the N-terminus by the addition of amino acid residues, wherein at least one of the amino acids provides an accessible sulfhydryl group. The accessible sulfhydryl group or groups may be utilized during conjugation or remain available for further conjugation. The term modified annexin includes annexin multimers.

Annexin Multimer: A combination of two or more monomeric modified annexin molecules of which the components of the multimer may be native or recombinant, or in any combination thereof; resulting in similar or improved membrane binding affinity over the monomeric annexin. A multimer composed of up to about 20 modified annexins is useful for the present invention. The preferred multimer composition is between 2 and about 10 modified annexins. One example of an annexin multimer is an annexin dimer, which can be composed of two modified annexins linked by disulfide bonds between accessible sulfhydryl groups on the modified annexins. The annexin dimer can be produced directly as a fusion protein using known expression systems, wherein the two annexin molecules can be connected by a peptide linker through

the accessible sulfhydryl groups. A dimeric molecule could contain additional functional sites, such as an endogenous radiolabel chelation site or an accessible sulfhydryl group for the attachment of one or more hexose residues, for example. Also, as defined herein, annexin multimer is covered by the term annexin.

Extension: A series of amino acids added to the N-terminus of the annexin molecule which provides a sulfhydryl group within ten amino acids from the N-terminus. Preferably the sulfhydryl group is within six amino acids of the N-terminus. More preferably the sulfhydryl group is the second residue in the N-terminal extension of six amino acids. The sulfhydryl group is preferably provided by a cysteine residue. This addition of amino acids provides a highly flexible means to attach functional moieties to the annexin terminal region.

$N_xS_y$  Chelating Compounds: As defined herein, the term " $N_xS_y$  chelating compound" includes bifunctional chelators that are capable of (i) coordinately binding a metal or radiometal and (ii) covalently attaching to an annexin molecule. Preferred  $N_xS_y$  chelating compounds have the  $N_2S_2$  (generally described in U.S. Patent No. 4,897,225 or 5,164,176 or 5,120,526),  $N_3S$  (generally described in U.S. Patent No. 4,965,392),  $N_2S_3$  (generally described in U.S. Patent No. 4,988,496),  $N_2S_4$  (generally described in U.S. Patent No. 4,988,496),  $N_3S_3$  (generally described in U.S. Patent No. 5,075,099) or  $N_4$  (generally described in U.S. Patent No. 4,963,688 and 5,227,474)

cores. Particularly preferred  $N_xS_y$  chelating compounds have  $N_2S_2$  and  $N_3S$  cores. Exemplary  $N_xS_y$  chelating compounds are described in Fritzberg et al., Proc. Natl. Acad. Sci. USA 85:4024-29, 1988; in Weber et al., Bioconj. Chem. 1:431-37, 1990; and in the references cited therein, for instance. For the purpose of this description, the prototypical  $N_xS_y$  chelating compound is an  $N_2S_2$  chelating compound. A chelating compound which complexes a metal or radiometal is termed a "chelate".

Esterase-sensitive Chelating Compound: Amide thiolate acetate ester chelating compounds of the  $N_xS_y$  family wherein the esterase-sensitive chelating compound connects the radionuclide metal to the annexin. As defined herein, the term "esterase-sensitive chelating compounds" includes those described in U.S. Patent No. 5,112,953 and 5,175,257. The esterase-sensitive chelating compounds is cleaved during the metabolism of the radiolabeled protein in the liver to generate radioactivity associated catabolites of favorable redistribution, resulting in the reduction of background radioactivity adjacent to the heart and lungs. A preferred esterase-sensitive chelating compound is  $N_3S$ -serylsuccinate as described in U.S. Patent No. 5,112,953.

Radiolabeled Annexin: An annexin having a radionuclide complexed therein.

Radiolabeled Annexin-Galactose: A galactose-derivatized annexin having a radionuclide complexed therein.

Hexose moiety: a composition of one or more six carbon sugar (hexose) residues, or derivatives based upon such moieties recognized by Ashwell receptors or other liver receptors, such as the mannose/N-acetylglucosamine receptor, which are associated with endothelial cells and/or Kupffer cells of the liver, or by the mannose 6-phosphate receptor. A hexose moiety includes a hexose cluster. A hexose moiety also includes more than one hexose moiety conjugated at separate sites to another molecule such as annexin.

Hexose Cluster: A construct having a plurality of hexose residues (including derivatives) configured to be recognized by a liver receptor. Such clusters are preferably constructed of hexose residues connected in a branched configuration (via a linking moiety, which in turn is connected to a branching structure), and are attached to annexin or to other components of a hexose cluster containing conjugate via a single point of attachment (through a linking moiety). Preferably, the branching network consists of two or three pronged branches, i.e., consists of 2, 4, 8, 16, 32 or 64 hexose residues or consists of 3, 9, 27, or 81 hexose residues.

Galactose Cluster: A construct having from about 3 to about 64 galactose residues connected in a branched configuration. Preferably, the branching network consists of two or three pronged branches, i.e., consists of 2, 4, 8, 16, 32 or 64 galactose residues or consists of 3, 9, 27, or 81 galactose residues.

Radiolabeled Annexin-Galactose Cluster: A galactose cluster-derivatized annexin having a radionuclide complexed therein.

5      Conjugate: A conjugate encompasses chemical conjugates (covalently or non-covalently bound), fusion proteins and the like.

10      Cleavable Linker: A bifunctional linker recognized by liver enzymes in such a way that the linker is enzymatically cleavable to produce a hydrophilic catabolite. In some preferred embodiments of the present invention, the cleavable linker connects the chelating compound to the hexose moiety. The cleavable linker functions for enhancement of renal excretion and hepatobiliary excretion so as to increase target to background ratio enabling early detection of atrial and venous thrombi. Examples of this type of linker includes a polymer, such as monosaccharides, polysaccharides, polyamino acids, hydroxyalkyl acrylamides, polyethylene glycol based hydrophilic polymers, biodegradable polymers containing an ether or ester linkage, as well as dextran and hemisuccinyl esters.

20  
25      It will be evident to one of ordinary skill in the art that the references throughout the application to annexin includes modified annexin, as well as annexin multimer, unless otherwise specified or apparent from the context. One of ordinary skill in the art will also understand that, although galactose is described below as a representative example, any reference to galactose

includes other hexose moieties, such as N-acetylgalactosamine and clusters thereof.

5 The present invention is directed to annexin-hexose containing conjugates, and radiolabeled annexins, including radiolabeled annexin-galactose cluster  
10 conjugates, and the use thereof for diagnostic imaging purposes. Radiolabeled annexins of the present invention are characterized by the following: rapid accretion to target cell sites characterized by anionic phospholipids; short circulating half-life; in vivo stability against metabolic degradation or radionuclide separation from chelate; and amenability to packaging in cold kit format.

15 Radiolabeled annexin-galactose conjugates also exhibit the aforementioned characteristics, albeit to different degrees. For example, radiolabeled annexin-galactose conjugates exhibit a shorter circulating half-life than their radiolabeled annexin counterparts. Also, radiolabeled annexin-galactose conjugates generally exhibit a lower binding affinity for target sites than  
20 their radiolabeled annexin counterparts. Successful diagnostic imaging involves both target site accumulation of signal and rapid elimination of non-targeted signal. Consequently, the enhanced elimination from a recipient's circulation of the radiolabeled annexin galactose  
25 conjugates provides a distinct opportunity to achieve diagnostic images in a shorter time frame.

Moreover, target site signal diminishes over time as a result of radioactive decay. In addition, metabolism

of the target associated material also diminishes target signal. Consequently, shorter time frame imaging enhances the target:non-target ratio and total target signal, thereby garnering improved diagnostic information.

Radiolabeled annexin-galactose cluster conjugates of the present invention combine desirable features of radiolabeled annexins and radiolabeled annexin-galactose conjugates. For example, radiolabeled annexin-galactose cluster conjugates exhibit a shorter circulating half-life than their radiolabeled annexin counterparts. Also, radiolabeled annexin-galactose cluster conjugates generally exhibit a higher binding affinity for target sites than their radiolabeled annexin-galactose conjugate counterparts. Consequently, the enhanced elimination from a recipient's circulation and the substantial binding affinity maintenance of the radiolabeled-annexin-galactose cluster conjugates of the present invention offer the opportunity to achieve clear diagnostic images in a shorter time frame.

An embodiment of the present invention is directed to annexin-containing conjugates suitable for radiolabeling with a diagnostic imaging agent including:

an annexin;

a hexose moiety; and

an  $N_xS_y$  chelating compound associated with the annexin.

Radiolabeled annexin-galactose cluster conjugates

suitable for imaging vascular thrombi anywhere in the recipient (but particularly in or near the heart) are also contemplated, which radiolabeled annexin-galactose cluster conjugates incorporate an annexin, a cluster of galactose residues, an  $N_xS_y$  chelating compound, and further a diagnostic radionuclide complexed by the chelating compound.

A preferred embodiment of the present invention involves annexin-containing conjugates suitable for radiolabeling with a diagnostic imaging agent including:

an annexin;

a cluster of galactose residues; and

an  $N_2S_2$  chelating compound associated with the annexin.

Radiolabeled annexin-galactose cluster conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled annexin-galactose conjugates incorporate an annexin, a cluster of galactose residues, an  $N_2S_2$  chelating compound, and further a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention includes:

a modified annexin, wherein the modification provides an accessible sulfhydryl group; and

a hexose moiety recognized by a mammalian liver receptor, wherein the hexose moiety is conjugated to the annexin.

Radiolabeled modified annexin-hexose moiety conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled annexin-hexose moiety conjugates incorporate a modified annexin, a hexose moiety, and further a diagnostic radionuclide conjugated to the modified annexin.

Another preferred conjugate of the present invention includes:

a modified annexin, wherein the modification provides an accessible sulfhydryl group;

a hexose moiety recognized by a mammalian liver receptor; and

a  $N_xS_y$  chelating compound, wherein the hexose moiety is conjugated to the modified annexin directly or via the chelating compound is conjugated to the modified annexin directly or via the hexose moiety.

Radiolabeled modified annexin-hexose moiety conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled modified annexin-hexose moiety conjugates incorporate a modified annexin, a hexose moiety, an  $N_xS_y$  chelating compound and further a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention includes:

an annexin multimer;

a hexose moiety recognized by a mammalian liver receptor; and

an  $N_xS_y$  chelating compound, wherein the hexose moiety is conjugated to the multimer directly or via the chelating compound and the chelating compound is conjugated to the multimer directly or via the hexose moiety.

5 Radiolabeled annexin multimer conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled annexin multimer conjugates incorporate an annexin multimer, a hexose moiety, an  $N_xS_y$  chelating compound, and further a diagnostic radionuclide complexed by the chelating compound.

Another preferred annexin conjugate of the present invention is those suitable for radiolabeling with a diagnostic imaging agent including:

an annexin; and

15 an esterase-sensitive  $N_xS_y$  chelating compound conjugated to the annexin.

Radiolabeled annexin conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled annexin conjugates incorporate an annexin, an esterase-sensitive  $N_xS_y$  chelating compound and further a hexose moiety and a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention includes:

25 a modified annexin, wherein the modification provides accessible sulfhydryl group; and

a  $N_xS_y$  chelating compound conjugated to the annexin. Radiolabeled modified annexin conjugates suitable for

imaging vascular thrombi are also contemplated, which radiolabeled modified annexin conjugates incorporate a modified annexin, an  $N_xS_y$  chelating compound and further a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention includes:

a modified annexin, wherein the modification provides an accessible sulfhydryl group; and

an esterase-sensitive  $N_xS_y$  chelating compound conjugated to the annexin.

Radiolabeled annexin conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled annexin conjugates incorporate an annexin, an esterase-sensitive  $N_xS_y$  chelating compound, and further a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention includes:

an annexin multimer; and

a  $N_xS_y$  chelating compound conjugated to the annexin.

Radiolabeled annexin multimer conjugates suitable for imaging vascular thrombi are also contemplated, which radiolabeled annexin multimer conjugates incorporate an annexin multimer, an  $N_xS_y$  chelating compound, and further a diagnostic radionuclide complexed by the chelating compound.

Another preferred conjugate of the present invention

includes:

an annexin multimer; and  
 an esterase-sensitive  $N_xS_y$  chelating compound  
 conjugated to the annexin.

5 Radiolabeled annexin multimer conjugates suitable for  
 imaging vascular thrombi are also contemplated, which  
 radiolabeled annexin multimer conjugates incorporate an  
 annexin multimer,

10 an esterase-sensitive  $N_xS_y$  chelating compound, and  
 further a diagnostic radionuclide complexed by the  
 chelating compound.

15 It will be evident to one of ordinary skill in the  
 art, that where a conjugate of the present invention has  
 three components (e.g., an annexin, a hexose moiety, and  
 chelating compound), a component may be bonded to one  
 another in a variety of orientations. For example, the  
 hexose moiety and a chelating compound may be bonded at  
 different sites on an annexin, or bonded to one another  
 and only one is bonded to the annexin directly.

20 For the visualization of vascular thrombi associated  
 with a number of pathological conditions, a radiolabeled  
 annexin such as a conjugate of an annexin with a  
 chelating compound complexed with an imaging  
 radionuclide, such as Tc-99m for example, is administered  
 25 to a recipient for whom such a diagnosis is desired. The  
 radiolabeled annexin or annexin portion of the  
 radiolabeled conjugate localizes rapidly to target sites  
 characterized by negatively charged surface

phospholipids, such as vascular thrombi. The radionuclide is selected for its ability to be visualized via one of the various techniques therefor, e.g., gamma camera imaging. Because of the rapid accretion of annexins to the target site and the short serum half-life (generally less than 30 minutes) of annexins (which is not significantly lengthened upon radiolabeling), imaging of those target sites proceeds with little exposure of non-target sites to radioactivity.

Diagnostic imaging is dependent on signal-to-noise ratio. Improvements involving either target signal accumulation or noise reduction will enhance the efficacy of the diagnostic imaging product. For targeted imaging, noise reduction is synonymous with reducing background radioactivity, particularly blood pool activity. Annexins, including annexin V, are rapidly removed by the liver; however, only a fraction of the activity is removed with each pass of the circulating conjugate through the liver.

To more efficiently remove the background activity from the blood, an improvement in liver extraction of annexin-containing conjugates could be employed. A preferred method to enhance liver extraction involves derivatizing the annexin-containing conjugate with a hexose moiety, such as galactose, that is recognized by a liver receptor. The efficiency of liver receptor extraction of the moiety recognized thereby will result in increased "per pass" removal by the liver of

derivatized annexin-containing conjugate (e.g., annexin-galactose conjugate) in comparison to the amount of underivatized annexin-containing conjugate so removed.

Further improvement in signal-to-noise ratio can be achieved by derivatizing radiolabeled annexin or an annexin-containing radiolabeled conjugate with a cluster of galactose molecules recognized by a liver receptor. The efficiency of liver receptor extraction of the moiety recognized thereby will result in increased "per pass" removal by the liver of derivatized annexin-containing conjugate (e.g., annexin-galactose cluster conjugate) in comparison to the amount of underivatized annexin-containing conjugate so removed. In addition, the multiple galactose residues arranged in a cluster are bound to the annexin conjugate component via a single point of attachment. Consequently, less or no reduction in annexin target binding affinity resulting from galactose cluster derivatization may be achieved in comparison to annexin-galactose conjugates.

Annexins are generally (with the most notable exception being annexin II) single chain, non-glycosylated proteins of approximately 33-72 kilodalton molecular weight. Annexins possess a number of biological activities associated with calcium ion-mediated binding.

Investigations have shown that annexins bind with high affinity to anionic membrane lipids in the presence of millimolar concentrations of calcium. In the presence

of calcium, these proteins have an especially high affinity for negatively charged phospholipids, such as phosphatidylserine, phosphatidylglycerol, phosphatidic acid, or phosphatidylinositol. See, for example, Funakoshi et al., Biochem. 26: 5572-78, 1987; and Tait et al., Biochem. 27: 6268-76, 1988. Such negatively charged phospholipids are associated with vascular thrombi (e.g., are located on the surface of activated human platelets).

Annexins exert anti-coagulatory effects. Coagulation inhibition is mediated by the binding of annexins to negatively charged surface phospholipids (e.g., present on the surface of activated platelets). This binding is believed to block the activation of clotting factors by such negatively charged surface phospholipids. Annexins localize to target sites bearing anionic phospholipids rapidly, i.e., in a matter of approximately 5 to 30 minutes depending on circulating levels thereof, but remain circulating in the serum for a somewhat longer time period (circulating half-life < 30 minutes).

Example III below discusses results of imaging experiments wherein vascular thrombi were visualized in planar images at an average time (following annexin administration) of 82 minutes.

Because of these properties, annexins or annexins conjugated to diagnostic or therapeutic agents may be employed in protocols for the *in vivo* diagnosis or treatment of vascular thrombi associated with a number of indications, such as DVT (deep vein thrombosis), PE

(pulmonary embolism), myocardial infarction, atrial fibrillation, problems with prosthetic cardiovascular materials, stroke and the like. Other indications associated with accumulation of activated platelets, for which the annexin conjugates of the present invention are useful, include the following: abscess imaging, restenosis post balloon angioplasty (PCTA), inflammation of joints (i.e., Rheumatoid arthritis), damaged endothelial cells (i.e., Alzheimer's disease), imaging of clots in cerebral arteries, occlusions in peripheral arteries, atrial thrombosis imaging, and imaging of coronary and carotid artery thrombi.

It is also important to characterize platelet populations in clinical, *in vitro* diagnostic, and basic research disciplines. Of the cell surface markers currently available to characterize platelets, many are not cross-reactive between species and may recognize all platelets as opposed to just the activated platelet population. It is believed that annexin selectively binds to activated platelets in many species. Thus annexin conjugates of the invention can be utilized as an alternative cell marker in research and diagnostics disciplines such as immunohistochemistry and flow cytometry. For example, conjugates of the present invention can be used to detect activated platelets in fixed tissues/tumors, blood smears, in animals with coagulopathies, and *in situ* in platelet activation assays in response to various chemical or infectious stimuli as

well as to detect activated platelets in blood, cell culture assays and platelet response assays. It will be appreciated by one of ordinary skill in the art that the conjugates of the present invention are useful for any purpose or indication, in which the targeting or binding of activated platelets would be desirable. Exemplary diagnostic protocols and experimentation employing radiolabeled annexins are set forth below to further elucidate this aspect of the present invention.

An example of a preferred annexin useful in the practice of the present invention is Annexin V, which was isolated by Bohn in 1979 from human placenta, a rich source of annexins, and termed Placenta Protein 4 (PP4). Annexin V has been expressed in E. coli, Iwasaki et al., J. Biochem., Vol 102, No.5, 1261-1273 (1987). Also, a full length cDNA clone of annexin V has been obtained and subcloned in expression vectors, thereby facilitating the production of fusion proteins containing annexin V (see generally, Tait et al., J. Biol. Chem., Vol. 270, No. 37, pp 21594-21599 (1995). Annexin V consists of four domains (four tandem, imperfect repeats of about 75 amino acid residues, Funakoshi et al., Biochem., 26: 8087-92, 1987), wherein each domain is made up of 5 alpha helices. From the side, the annexin V molecule appears crown-like with at least four calcium binding sites on its convex surface, through which annexin-phospholipid interactions are mediated. Other annexin molecules are also useful in the practice of the present invention, and the

discussions relating to annexin V herein apply generally to annexin molecules.

Because annexin V has a plurality of calcium binding sites, and because annexin V binding to negatively charged phospholipids is mediated by calcium, an engineered molecule consisting of one or more individual annexin V domains may be employed in imaging protocols of the present invention. Also, the annexin molecule may be partitioned at a position or positions different from the domain boundaries to provide an engineered molecule capable of calcium-mediated binding of anionic phospholipids. Also, annexin V may be altered at one or more amino acid residues, so long as the affinity of annexin V for anionic phospholipids is not significantly impaired. For example, the cysteine (position 316) amino acid residue of annexin V can be either deleted or replaced with alanine or other non-sulfur containing amino acids known to those of ordinary skill in the art; wherein, upon labeling, a monomerically radiolabeled annexin V is produced.

In another example, a native annexin can be modified at the N-terminus by adding amino acid residues to provide for an accessible sulfhydryl group or groups. This can be accomplished with the addition of at least a single cysteine residue near the N-terminus. The accessible sulfhydryl groups may be available after initial conjugation or is for further conjugation depending on the embodiment of the invention utilized.

A preferred modified annexin molecule, as defined herein, is a monomeric form of annexin V with an N-terminal extension of an amino acid sequence, wherein the sequence is selected so that it contains amino acids adjacent to an amino acid countacy an accesible sulfhydryl group cysteine. The selected sequence will improve target to normal organ ratios and superior clot imaging potential. The use of hydrophilic amino acids, in the sequence such as serine, glycine, theonine, aspartate, glutamate and others facilitates renal excretion of chelate products of catabolism following uptake of the radiolabeled annexin in normal organs, such as the liver. One preferred amino acid sequence added to at the N-terminus of annexin V is Ala-Cys-Asp-His-Ser-Met. An advantage of this particular configuration is that with the cysteine near the N-terminus, the sulfhydryl and neighboring amide groups provide for stable chelation of the technetium resulting in an  $N_3S$ -like stable chelate, as in endogenous radiolabeling of the annexin molecule.

Other model peptides that vary the sequence at the N-terminus of modified annexin V (for example, the sequence of George et al. in Proc. Natl. Acad. Sci. USA, 92:8358-62, 1995) can be prepared. These peptides are subjected to the same conditions of *in vitro* labeling with Tc-99m as described above. Peptides with suitable labeling characteristics can be chosen, and the sequence can be engineered into the recombinant annexin V molecule.

Success is indicated by preparation of a derivative with specific activity as high or higher than the current derivative as reported by Stratton et al, in Circulation, 92: 3113-3121, 1995; that is stable in vitro and in vivo; that retains membrane and thrombus binding activity; and that is not adversely altered with respect to blood clearance or biodistribution. If the above-identified criteria is met, the modified Annexin V molecule or dimer of the modified annexin V molecule is used according to the invention.

An annexin multimer composed of two or more modified annexin molecules which are limited by disulfide bonds between one or more of the accessible sulfhydryl groups on the respective annexins. A preferred annexin multimer, is an annexin dimer. A prototypical annexin multimer is an annexin dimer. The dimer is intended to have a higher affinity for the platelet membrane resulting from a slower dissociation rate. The dimer can then be accreted to a higher level or retained longer in the thrombus *in vivo*, thus improving the target-to-background ratio at any given time. The annexin multimer, as well as the annexin dimer can be produced as a fusion protein using known expression systems with the modified annexins connected by a peptide linker through the accessible sulfhydryl groups. A multimeric molecule can also contain additional functional sites, such as an endogenous radiolabeled chelation site or a free sulfhydryl to allow for attachment of a hexose moiety.

Where an annexin dimer is produced and includes a hexose moiety and a chelating compound, this dimer can then be used for generating stabilized monomers for further radiolabeling procedures via reduction of the disulfide bonds of the dimer.

The degree of annexin binding to phospholipids may be quantified by fluorescence quenching as described by Tait et al., J. Biol. Chem., 264: 7944-49, 1989.

Among annexins, annexin V has the strongest binding affinity ( $K_d < 10^{-10}$  M) for phospholipid vesicles containing 80% phosphatidylcholine and 20% phosphatidylserine under conditions comparable to plasma and extracellular fluid (1.2 mM ionized calcium, 0.15 M ionic strength). This binding is reversible and calcium dependent.

To decrease the background radiolabel activity, annexins (including modified annexins) may be derivatized with hexose moieties or hexose-based moieties. More specifically, annexins may be derivatized to incorporate one or more hexoses (e.g. six carbon sugar moieties) recognized by Ashwell receptors or other liver receptors, such as the mannose/N-acetylglucosamine receptor, which are associated with endothelial cells and/or Kupffer cells of the liver, or by the mannose 6-phosphate receptor. Exemplary of such hexoses are galactose, mannose, mannose 6-phosphate, N-acetylglucosamine, pentamannosyl phosphate, and the like. Other moieties recognized by Ashwell receptors, including glucose, N-

galactosamine, N-acetylgalactosamine, thioglycosides of galactose and, generally, D-galactosides and glucosides and the like, may also be used in the practice of the present invention. Thus, hexose moieties useful in the practice of the present invention include a variety of galactose, mannose and glucose sugars recognized by liver receptors. Preferred hexose moieties are hexose sugars, such as galactose and galactose clusters. It should be understood that galactose and galactose clusters are encompassed by the term hexose moiety. It should also be recognized that the use of the term galactose in the discussion is understood to encompass not only galactose, but N-galactosamine, N-acetylgalactosamine, thioglycosides of galactose and generally, D-galactosides as well; the term mannose is understood to encompass mannose, mannose-6-phosphate, pentamannosyl phosphate, and the like; the term glucose, is understood to encompass glucose, glucosides, and the like. For purposes of the invention the most preferred galactose is N-acetylgalactosamine.

Galactose is the prototypical hexose employed for the purposes of this description. Galactose thioglycoside conjugation to a protein is preferably accomplished in accordance with the teachings of Lee et al., "2-Imino-2-methoxyethyl 1-Thioglycosides: New Reagents for Attaching Sugars to Proteins," Biochemistry, 15(18): 3956, 1976. Another useful galactose thioglycoside conjugation method is set forth in Drantz et al,

5

20

25

Hexose clusters are preferably employed in the practice of the present invention. Galactose clusters are the prototypical hexose clusters employed for the purposes of this description. A cluster containing the

galactose, N-aceylgalactosamine, is a preferred galactose clusters of the present invention. Design of hexose clusters of the present invention is conducted with the following criteria in mind, as set forth in the context of the design of a galactose cluster:

- 1) Number of Galactoses in a Cluster;
- 2) Distance Between Galactoses in the Cluster; and
- 3) Distance Between Galactose Cluster and the Annexin Conjugate Component.

With regard to criteria number 1, literature indicates that galactose receptors on the surface of human hepatocytes are grouped as heterotrimers and, perhaps, bis-heterotrimers. See, for example, Hardy et al., Biochemistry, 24: 22-8, 1985. For optimal affinity to such receptors, it is contemplated within the present invention that each galactose cluster should preferably contain at least three galactose residues. In general, the greater the number of galactose residues in a cluster, the greater the propensity for the cluster to be recognized by liver receptors.

Increased galactose cluster size may impair annexin binding to target. If significant impairment in annexin binding to target (e.g. reduction to < 20% of native annexin binding) is observed, a longer linker between the two moieties may be used or such large clusters should not be employed in radiolabeled annexin-galactose cluster conjugates of the present invention.

With respect to criteria number 2, the galactose



receptors within each trimer are separated from each other by distances of 15, 22 and 25 angstroms. Consequently, it is contemplated within the present invention that the galactoses within a cluster should preferably be separated by flexible linkers allowing separation of at least 25 angstroms. The spacing between galactose residues is likely to be more important if the number of galactose residues is small. With larger constructs, appropriate spacing is likely to occur with respect to galactose residues that are not immediate neighbors (i.e., sugar residues that are farther apart than those that are immediate neighbor). Assuming an average bond length of 1.5 angstroms, preferred galactose clusters of the present invention are characterized by separation of neighboring galactose residues by about 10 bond lengths or more. Other preferred constructs involve galactose clusters characterized by separation of neighboring sugar residues by about 25 bond lengths or more.




Regarding criteria number 3, the distance between the annexin and the galactose cluster should be sufficient to obviate any adverse steric effects upon annexin binding to target caused by the size or orientation of the galactose cluster. This distance is preferably greater than about 7 bond lengths or about 10 angstroms. If necessary, an extender molecule is incorporated between the galactose cluster and the linker (which joins the galactose cluster and the annexin component) or between

the annexin and the linker to provide the requisite distance.

While the foregoing parameters appear to be optimal for galactose, it should be noted that these factors may vary with other hexoses or mixtures thereof, which may or may not bind to the same receptors, or may bind differently. Given the teachings in this application, one of ordinary skill in the art can, using available synthesis techniques, attach an annexin and an active agent to other hexose clusters and identify those constructs which provide optimal performance.

Any branched sugar structures that meet the criteria described above may be employed in the practice of the present invention. Preferred galactose clusters of the present invention are of the following structures:








wherein X is preferably H or methyl, resulting in galactose clusters bearing 4, 8, 16 and 32 galactose residues, respectively. Further iteration in the branching scheme allows expansion of the galactose cluster to include 32, 64, etc. galactose residues. In addition, the linker moiety between the hexose itself and the branching structure (shown as  $-\text{S}-(\text{CH}_2)_4-\text{NX}-$ ) may be variable in length.

Alternative branching structures may also be employed in the design of galactose clusters in accordance with the present invention. For example, other constructs wherein the branching results in a doubling of the number of galactose residues may be employed. In addition,

constructs wherein branching results in a tripling or other convenient multiplying of the number of galactose residues are also contemplated by the present invention.

Another potential branching construction is based upon the molecule bis-homotris:  $(\text{HO-CH}_2)_3\text{-C-NH}_2$ . The sulphydryl-containing derivative of this molecule may also be used. In this embodiment of the present invention, each arm of the bis-homotris molecule is extended and terminated in a carboxylic acid:  $(\text{HO}_2\text{C-(CH}_2)_y\text{-Z-(CH}_2)_3\text{-C-NH}_2)$ , where Z is S or O and y ranges from 1 to about 10. For this embodiment of the present invention, a preferred galactose cluster is characterized by the following structures:



wherein X is preferably H or methyl; y ranges from 1 to about 10; and Z is O or S. The above structures bear 3, 9 and 27 galactose residues, respectively. Further iteration of the branching allows expansion to include

81, etc. galactose residues.

Also, X may be a lower alkyl moiety (composed of two to twelve carbons) different from methyl, such as ethyl, t-butyl and the like. X may also be a lower alkyl group bearing a heteroatom, such as a lower alkyl acid, ester, aldehyde, ketone or ether.

In embodiments of the present invention wherein the chelating compound is characterized by a single functional group available and suitable for conjugation, the annexin, chelating compound and galactose cluster components are preferably joined via a trifunctional linker. Functional groups that are "available" for conjugation are those that are not prevented by steric constraints from conjugate formation. Functional groups that are "suitable" for conjugation are those that are capable, in a chemical sense, of reacting with available functional groups associated with other conjugate components. In addition, conjugation of "suitable" functional groups does not substantially impair a necessary function of the component with which the functional group is associated. For example, a functional group located in the complementarity determining region of an antibody targeting moiety will generally not be "suitable" for conjugation, because the targeting ability of the antibody is likely to be substantially impaired by such binding.

Useful trifunctional linkers are amenable to binding with functional groups available on the three conjugate

components or any extender moieties employed in conjugate construction. A useful trifunctional linker is lysine, wherein the alpha-amino, epsilon-amino and carboxyl functional groups are used. One of ordinary skill in art the could identify other trifunctional linkers and use the same in the practice of the present invention as set forth herein.

The following variations for the order of component (specifically the modified annexin, hexose moiety and chelating compound) conjugation, are contemplated within the present invention:

the hexose moiety conjugated to the modified annexin, through an accessible sulfhydryl group or groups, and the chelating compound conjugated to the modified annexin through a lysine residue on the modified annexin; a chelating compound conjugated to the modified annexin, through accessible sulfhydryl group or groups, and the hexose moiety conjugated to the modified annexin through a lysine residue on the modified annexin; chelating compound conjugated to hexose moiety, which in turn is conjugated to the modified annexin through accessible sulfhydryl group or groups; the hexose moiety conjugated to the chelating compound, which in turn is conjugated to the modified annexin through accessible sulfhydryl groups; chelating compound conjugated to the hexose moiety which in turn is conjugated to the modified annexin through the lysine residues of the modified

annexin, thus leaving the accessible sulfhydryl groups available for further conjugation; hexose moiety conjugated to chelating compound which in turn is conjugated to the modified annexin through lysine residues, again leaving the accessible sulfhydryl groups available for further conjugation; and other like variations.


Cleavable linkers may be employed in the present invention. Such linkers are bifunctional moieties capable of binding with the trifunctional linker in such a way as to connect the chelating compound to the trifunctional linker, which in turn is connected to the annexin component and cluster component. Examples of suitable cleavable linkers include monosaccharides, polysaccharides, polyamino acids, hydroxyalkyl acrylamides (e.g. HPMA), polyethylene glycol based hydrophilic polymers, biodegradable polymers which contain an ether or ester linkage, as well as dextran and hemisuccinyl esters, and the like.

Extender molecules useful in the present invention are bifunctional moieties capable of binding with either the annexin component and the linker or the galactose cluster component and the linker. Suitable extender molecules include aminocaproate moieties,  $\text{HS}-(\text{CH}_2)_n\text{COOH}$  or an activated ester form thereof wherein  $n$  ranges from 2 to about 5, and the like. One of ordinary skill in the art is capable of identifying and using other suitable extender molecules as described herein. Alternatively,

the extender function can be served by an appropriately constructed linker.

Also, binding facilitation moieties may also be employed in the present invention. Such moieties are bifunctional and facilitate binding the conjugate components, e.g., galactose cluster, annexin, chelating compound, linker, and extender. Examples of such binding facilitation moieties include urea functionalities, thiourea functionalities, succinate bridges, maleimides and the like. Such binding facilitation moieties are amenable to identification and use by those of ordinary skill in the art.

An example of a linker-extender-binder facilitation system is shown below:



wherein the alpha-amine of the lysine linker is bound via a urea or thiourea functionality to an amino caproate spacer (which, in turn, binds to a galactose cluster that is not shown); the lysine carboxylate is available for linkage to a chelating compound (not shown); and the epsilon-amine of the lysine linker is available for linkage to a lysine residue of the annexin component (not shown) via a succinate bridge. Other amino acid residues of the annexin component, such as cysteine, may also be employed for binding purposes. Alternatively a maleimide-S-(CH<sub>2</sub>)<sub>n</sub>CO- binding facilitation moiety-

extender combination may be employed to link the sugar residue with the lysine. An example of a conjugate joined by a trifunctional linker is discussed in Example VII.

5           When the chelating compound component of the conjugate is characterized by greater than one functional group available and suitable for conjugation, the galactose cluster may be linked to the chelating compound component which, in turn, is linked to the annexin component of the conjugate via two or more bifunctional linkers. Alternatively, the hexose cluster, such as N-acetylgalactosamine, is linked to the accessible sulfhydryl group of the modified annexin and the chelating compound component is linked to the annexin component via the lysine residue on the annexin. In another possible configuration, the chelating compound is linked to a hexose cluster through a trifunctional linker, wherein the hexose cluster conjugated to the chelating compound is then linked to the accessible sulfhydryl group of the modified annexin through a binding facilitation moiety. The above possible configuration can also include a cleavable linker between the chelating compound and the linker attaching the cluster to the annexin. Preferably, the annexin component of the conjugate is attached last in the formation of a galactose cluster-chelating compound-annexin conjugate. Suitable bifunctional linkers and linking methodologies can be identified and employed by

one of ordinary skill in the art. An example of such a chelating compound and such a conjugation methodology is set forth in Example V.

5 Annexin V modified with a hexose moiety, such as a galactose cluster containing, for example, a N-acetylgalactosamine sugar, is intended to clear faster from the blood by the liver, allowing one to achieve optimal thrombus-to-background ratios soon after  
10 administration of radiolabeled agent. Attachment of the cluster near the N-terminus of the modified annexin intended to not affect the binding of the annexin moiety to thrombi.

15 Annexin conjugation to the hexose moiety and the chelating compound via chemical methods can occur either prior to (post-formed approach) or following (pre-formed approach) complexation of a radiometal with the chelating compound. In a preferred embodiment, a hexose moiety is attached to the annexin component prior to the addition of the radiolabel. The chemical conjugation is  
20 preferably conducted following radiometal complexation, unless the chelating compound employed in the conjugate is capable of binding the radionuclide with rapid kinetics at room temperature. Annexin V is radiolabeled with an imaging radionuclide for use in the present  
25 invention.

Radionuclides useful within the present invention include penetrating photon emitters including gamma-emitters and X-ray emitters. These rays accompany

nuclear transformation such as electron capture, beta emission and isomeric transition. Radionuclides useful include those with photons between 80 and 400 keV and positron producers, 511 keV annihilation photons and acceptable radiation dose due to absorbed photons, particles and half life. Radionuclides suitable for use in the present invention are known in the art and include  $^{18}\text{F}$ ,  $^{64}\text{Cu}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{100}\text{Pd}$ ,  $^{212}\text{Bi}$ ,  $^{212}\text{Pb}$ ,  $^{109}\text{Pd}$ ,  $^{67}\text{Cu}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{94}\text{Tc}$ ,  $^{95}\text{Ru}$ ,  $^{105}\text{Ru}$ ,  $^{99}\text{Rh}$ ,  $^{105}\text{Rh}$ ,  $^{111}\text{In}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{153}\text{Sm}$ ,  $^{177}\text{Lu}$ ,  $^{170}\text{Lu}$ ,  $^{189}\text{Pt}$ ,  $^{193}\text{Pt}$ ,  $^{199}\text{Au}$ ,  $^{197}\text{Hg}$  and the like.


Tc-99m is a preferred radionuclide for the practice of the present invention. Tc-99m has been stably bound to annexin V in accordance with the present invention at both low and high specific activity ( $0.53 \mu\text{Ci}/\mu\text{g}$  -  $101.2 \mu\text{Ci}/\mu\text{g}$ ). Adequate radiochemical yields and good radiochemical purities were obtained. Activated platelet binding studies were also conducted, and the radiolabeled annexin V conjugates bound to activated platelets well.

$\text{N}_x\text{S}_y$  chelating compounds include bifunctional chelating compounds that are capable of coordinately binding a metal or radiometal and covalently attaching to an annexin molecule. Preferred  $\text{N}_x\text{S}_y$  chelating compounds have the  $\text{N}_2\text{S}_2$  (generally described in U.S. Patent No. 4,897,225 or 5,164,176 or 5,120,526),  $\text{N}_3\text{S}$  (generally described in U.S. Patent No. 4,965,392),  $\text{N}_2\text{S}_3$  (generally described in U.S. Patent No. 4,988,496),  $\text{N}_2\text{S}_4$  (generally described in U.S. Patent No. 4,988,496),  $\text{N}_3\text{S}_3$  (generally described in U.S. Patent No. 4,988,496),  $\text{N}_3\text{S}_3$  (generally described in U.S. Patent No. 5,075,099) and  $\text{N}_4$  (generally


described in U.S. Patent No. 4,963,688 and 5,227,474) cores. Particular preferred  $N_xS_y$  chelating compounds have the  $N_2S_2$  and  $N_3S$  cores.  $N_2S_2$  and  $N_3S$  chelating compounds are known in the art. For example, preferred  $N_2S_2$  chelating compounds are described in U.S. Patent No. 4,897,225, and preferred  $N_3S$  chelating compounds are generally described in U.S. Patent No. 4,965,392 and U.S. Patent No. 5,112,953.

The  $N_2S_2$  chelating compounds are diamide, dimercaptide bifunctional chelators of the  $N_xS_y$  family capable of stably complexing a radionuclide through two nitrogen atoms and two sulfur atoms that are appropriately positioned.  $N_2S_2$  chelating compounds are generally described in U.S. Patent No. 4,897,225.

Preferred chelating compounds with an  $N_2S_2$  core also include diamine dimercaptide chelating compounds having the following biphenyl backbone, which are described in U.S. patent application Serial No. 08/463,232:



OR



wherein  $n = 0$  to  $1$  and one or more  $T$  substituents incorporate a functional group available and suitable for conjugation with another conjugate component. Such exemplary functional groups include hydrophilic groups that are amenable for renal excretion as opposed to hepatobiliary excretion.

The  $N_3S$  chelating compounds are triamide, mercaptide bifunctional chelators of the  $N_xS_y$  family capable of stably complexing a radionuclide through three nitrogen atoms and one sulfur atom that are appropriately positioned. Preferred  $N_3S$  chelating compounds are described in U.S. Patent No. 4,965,392 and 5,091,514.

The present invention applies this stable chelation technology to exploit the thrombus-targeting ability of annexin molecules, thereby providing an imaging agent which is able to rapidly visualize vascular thrombi in vivo. The radiolabeled annexins, including radiolabeled annexin-galactose conjugates and radiolabeled annexin-galactose cluster conjugates, of the present invention can be used to obtain thrombus images which reduce or eliminate the high level of background radioactivity that results from metabolically degraded radiolabeled conjugate. The radiolabeled annexins, including

radiolabeled annexin-galactose conjugates and radiolabeled annexin-galactose cluster conjugates, also avoid clinically unacceptable toxicity to non-target cell sites. Tc-99m radiolabeled annexin V performed better than I-123 in the pig studies set forth in Example III hereof.

Radiolabeling of annexin V with a radionuclide using an  $N_2S_2$  or  $N_3S$  chelating compound may be conducted using either a pre-formed or a post-formed approach. That is, the radionuclide is either complexed within the chelating compound prior to (pre-formed) or following (post-formed) conjugation of the chelating compound to annexin V. The pre-formed approach is preferred and suitable procedures therefor are set forth in Examples I, II, IV, and XV. Further studies indicated that the pre-formed radiolabeling approach yields two radiometric peaks by HPLC analysis. The appearance of two peaks results from cysteine-conjugated annexin V as well as lysine-conjugated annexin V. Thus, the cysteine amino acid of annexin V can be either deleted or replaced with other non-sulfur containing amino acid to result in a monomerically labeled annexin V, see Example IX. Using the preformed approach to radiolabel the modified annexin V also results in a monomerically labeled annexin V. The conjugation can occur through the accessible sulfhydryl group of a modified annexin, instead of through the lysine residue of the modified annexin.

For embodiments of the present invention wherein


10

15

An esterase-sensitive chelating compound can be used

in a variety of the embodiments of the present invention. Such embodiments include annexin-chelating compound conjugates, including modified annexin-chelating compound conjugates, as well as multimer-chelating compound conjugates. A hexose moiety can be utilized in in the above embodiments. An esterase-sensitive chelating compound is cleaved during the metabolism of the radiolabeled protein in the liver to generate radioactivity associated catabolites of favorable redistribution, resulting in the reduction of background radioactivity adjacent to the heart and lungs. It is desirable to have a greater fraction of the technetium-99m released rapidly by the liver, and then excreted by the kidneys. This is accomplished by conjugating an esterase-sensitive chelating compound to the annexin protein. The ester component of this chelating compound is cleaved during metabolism of labeled protein in the liver, resulting in greater release of radioactivity from the liver into the blood and excreted from the kidneys. A preferred esterase-sensitive chelating compound one containing N<sub>3</sub>S-serylsuccinate, as described in U.S. Patent No. 5,112,953.

The following structure may be employed with annexin to obtain the above mentioned results:



wherein R is any acid labile or base labile sulfur protecting group known in the art, such as ethoxyethyl and R<sup>1</sup> is activated esters which include N-hydroxysuccinimide tetrafluorophenyl and tetrafluorophenyl and tetrafluoro-thiophenyl esters.

One of ordinary skill in the art is versed in the nature and specificity of functional moieties, and such nature and specificity are also discussed herein.

As indicated previously, annexin molecules may be modified by the addition of from about 2 to about 6 terminal amino acid residues to facilitate the conjugation reaction between the annexin molecule and a chelating compound, or between the annexin molecule and a hexose moiety, or between the annexin molecule and either a linker or a hexose moiety. For example, terminal amino acid residues may be added to provide a sulfhydryl group or to provide a group capable of derivatization to a maleimide group, with such sulfhydryl and maleimide groups available for the conjugation reaction. This modification may be made by protein chemistry methods or

via production of an appropriate fusion protein or other techniques useful therefor.

5 In one embodiment of the invention, a recombinant annexin molecule is modified at the N-terminus by the addition of amino acid residues, provided that at least one of the amino acids provides an accessible sulfhydryl group. For example, this can be accomplished with the addition of at least a single cysteine residue near the N-terminus. A preferred modified annexin as defined herein is a monomeric form of annexin V with an N-terminal extension of the preferred amino acid sequence of Ala-Cys-Asp-His-Ser-Met. One advantage of this configuration is that with the cysteine near the N-terminus, the sulfhydryl and neighboring amide groups provide for chelation of the technetium resulting in an N<sub>3</sub>S-like stable chelating compound, for direct labeling. It is intended within the present invention that adjacent amino acid donor atoms from amino acid moieties will wrap around the base of the metal oxo bond in a square pyramidal arrangement, thus achieving stability, similiar to stable chelation achieved by utilizing an N<sub>3</sub>S chelating compound.

20 Another example of the production of a modified annexin molecule contains a single cysteine residue as the second residue in an N-terminal extension of six amino acids. In this example, the single natural internal cysteine residue has been mutated to a non-sulfur containing amino acid, such as alanine or serine.

A method for mutating the cysteine residue is provided by Example VIII. This amino acid extension provides a highly flexible means to attach functional moieties to the N-terminal region of an annexin, such as annexin V. The procedures described herein successfully produce monomeric and dimeric modified annexin V molecules. It is apparent that the proteins are substantially pure and of the expected molecular weight. In addition, any production of a modified annexin V dimer can be quantitatively converted to the monomer, by treatment with 2-mercaptoethanol, confirming the presence of the expected disulfide linkage.

Measurements of the membrane binding properties of modified annexin, and the modified annexin dimer in a competition assay against native monomeric annexin V have been performed. The dimer is 6-fold more potent than monomeric annexin V in displacing labeled ligand from the membrane. Pharmacokinetic studies disclosed herein have shown that the increased molecular weight of the modified annexin dimer does not slow its rate of disappearance from the blood. In fact the modified annexin V dimer is removed from the blood at about the same rate as wild-type monomeric annexin V. See Table I below.

Table I  
Pharmacokinetics of  $^{125}\text{I}$  Iodinated modified annexin V  
Dimer and Annexin V in Mice

Time Since Injection	Radioactivity in Blood (%ID/g)	
	Annexin V	Modified Annexin V Dimer
15 min	1.27 $\pm$ 0.12	1.70 $\pm$ 0.02
60 min	0.60 $\pm$ 0.04	0.36 $\pm$ 0.04
240 min	0.06 $\pm$ 0.01	0.06 $\pm$ 0.02

The disclosure of the present invention shows that the dimeric molecule of annexin V is likely to have a higher affinity for the platelet membrane, most likely due to a slower dissociation rate. Thus, this molecule may be taken up to a higher level or retained longer in the thrombus in vivo, thus improving the target-to-background ratio at any given time.

Radiolabeled annexins, including radiolabeled annexin-galactose conjugates and radiolabeled annexin-galactose cluster conjugates, of the present invention offer an additional advantage over the previously prepared I-123-labeled annexins, in that they are amenable to packaging in a cold kit. That is, the annexin or annexin-galactose or annexin-galactose cluster or galactose cluster and chelating compound components may be individually vialled and provided separately from the Tc-99m component (and, possibly, vialled separately

from each other). When a patient requiring a thrombus image is identified, a cold kit may be ordered or retrieved from storage; Tc-99m may be obtained from a radiopharmacy or other source thereof; the pre-formed or post-formed chelation/complexation process is performed; the radiolabeled annexin, radiolabeled annexin-galactose conjugate or radiolabeled annexin-galactose cluster conjugate is administered to the patient; and the patient is subsequently imaged. For radiolabeled annexin-galactose conjugates, an annexin-galactose conjugate is preferably prepared and vialled in the kit. For radiolabeled annexin-galactose cluster conjugates, a chelating compound-annexin-galactose cluster conjugate is preferably prepared and vialled in the kit, although annexin-galactose cluster/chelating compound two component or other multi-component kits may also be used.

It is contemplated by the present invention that Tc-99m can be directly complexed through a specific bioengineered site on annexin V, without the need for intermediate production of an amine-directed active ester containing Tc-99m. To facilitate direct labeling, the modified Annexin V N-terminal sequence can consist of Ala-Cys-Asp-His-Ser-...etc, for example. As the cysteine is near the N-terminus, the sulfhydryl and neighboring amide groups are able to chelate the technetium resulting in an endogenous MAG<sub>3</sub>-like, stable technetium chelate. A defined chelate system through an endogenous protein sequence can be achieved wherein the adjacent amide donor

atoms from amino acid residues wrap around the base of the metal oxo bond in a square pyramidal arrangement; thus, providing for stable chelation similar to stable chelation achieved when using an N<sub>3</sub>S chelate in a preformed labeling method. For the endogenous chelationm technetium-99m is the preferred radionuclide.

Lyophilization and vialing of the conjugate components in a sterile, pyrogen-free environment may be accomplished via techniques, known to persons skilled in the art of good manufacturing practices, particularly as such practices relate to biological materials.

Radiolabeled annexins, including radiolabeled annexin-galactose conjugates or radiolabeled annexin-galactose cluster conjugates, of the present invention are administered in such amounts as to deliver a diagnostically effective amount of radionuclide to target sites. Appropriate administered doses depend on a variety of factors that are largely patient specific, and practitioners in the art will consider such factors in the practice of the present invention. The components of the radiolabeled annexin also impact dose amounts in ways that are known to be or are routinely ascertainable by practitioners in the art. In general, radiolabeled annexin, including radiolabeled annexin-galactose conjugate or radiolabeled annexin-galactose cluster conjugate, is administered to large mammals at a dose ranging between about 0.3 and about 300 micrograms/kg body weight of the recipient, with from about 3 to about

10 micrograms/kg preferred, depending upon the physiological characteristics of the patient and the ailment involved or suspected.

5 A practitioner in the art is capable of identifying an appropriate dose and administration route for a given recipient with a given ailment.

10 Radiolabeled annexins, including radiolabeled annexin-galactose conjugates or radiolabeled annexin-galactose cluster conjugates, of the present invention may be administered in any convenient manner therefor. For example, intravenous infusion may be employed to administer radiolabeled annexins. Other routes of administration also find utility in the practice of the present invention. Exemplary additional administration routes are injection by the arterial (e.g., coronary artery), intracoronary, intralymphatic, intrathecal, or other intracavity routes, and the like.

15 After administration of the radionuclide, depending upon the nature of the radionuclide and the purpose of the administration, the recipient may be subject to various procedures for detection of radioactive emissions from the site or sites at which the radionuclide localizes. For example, conjugates containing Tc-99m are imageable by a gamma camera.

20 The invention is further described through presentation of the following examples. These examples are offered by way of illustration, and not by way of limitation.

Example I**Procedure for Radiolabeling an Annexin - N<sub>2</sub>S<sub>2</sub> Chelate Conjugate**

5           Annexin V can be isolated from a variety of tissue extracts, such as liver, lung and placenta, in accordance with procedures set forth in Funakoshi et al., Biochem. 26: 8087-92, 1987); Tait et al., Biochem. 27: 6268-76, 1988; and U.S. Patent No. 4,937,324, for example. In addition, annexin V can be expressed in E. coli, as described by Tait et al., Archives of Biochemistry and Biophysics 288: 141-44, 1991.

10           Annexin V was radiolabeled with Tc-99m by using a diamide dimercaptide N<sub>2</sub>S<sub>2</sub> chelating compound in accordance with the OncoTrac® (also referred to as Verluma™) Small Cell Lung Cancer Imaging Kit labeling procedure described in J. Nucl. Med. 32: 1445-51, 1991.

15           A preferred method for radiolabeling annexin V with Tc-99m constitutes a modified OncoTrac® kit procedure using C-18 Baker purified Tc-99m-N<sub>2</sub>S<sub>2</sub>-TFP. In this procedure, an acidified active ester solution was prepared by adding 0.16 ml of 0.2M hydrochloric acid: glacial acetic acid (14:2 ratio) to 0.6 ml of 2,3,5,6,-tetrafluorophenyl 4,5-bis-(S-1-ethoxy-ethyl-mercaptoacetamido)pentanoate (0.3 mg, 0.0005 mole freshly dissolved in 0.9 ml of isopropyl alcohol). Then 0.5 ml of this solution was added to 1.1 ml of Tc-99m-gluconate (prepared from 0.12 mg SnCl<sub>2</sub> · 2 H<sub>2</sub>O, 5.0 mg sodium

20

25

gluconate at pH 6.1-6.3, and 100 mCi/ml of [Tc-99m] pertechnetate, i.e., the first step in the OncoTrac® kit labeling procedure). The reaction mixture was heated at 75 °C for 15 minutes followed by cooling on ice. The resulting Tc-99m transchelated tetrafluorophenyl active ester derivative of Tc-99m-4,5-bis(thioacetamido) pentanoate was, optionally and preferably diluted with 2 ml water and purified by loading the reaction mixture on a conditioned C-18 cartridge (J.T. Baker, Phillipsberg, NJ), washing with 2.0 ml of water eight times followed by drying the column for 5 minutes, and eluting with 100% acetonitrile. The solvent was evaporated under a steady stream of N<sub>2</sub> gas. Then 0.15 ml of phosphate buffered saline (PBS), 0.15 ml of annexin V at 2.35 mg/ml, and 0.2 ml of 0.2 M bicarbonate (pH 10.0) were added for conjugation to the Tc-99m-N<sub>2</sub>S<sub>2</sub> chelate. After 20 minutes at room temperature, the Tc-99m-N<sub>2</sub>S<sub>2</sub>-annexin V conjugate was purified by passage through a G-25 SEPHADEX® (PD-10) column (available from Pharmacia, Piscataway, NJ) equilibrated with PBS. Fractions (1.0 ml) were collected, and those fractions containing annexin V were pooled. Protein concentration was determined by UV absorption at 280 nm. Tc-99m-annexin V (300 -350 µg) conjugate solution was diluted and stored in PBS containing bovine serum albumin (BSA) at a final concentration of 15-20 mg BSA/ml PBS prior to injection.

#### Example II

**Procedure for Radiolabeling an Annexin -  $N_3S$  Chelate  
Conjugate**

S-benzoylmercaptoacetylglycylglycylglycine (S-benzoyl  
MAG<sub>3</sub>) is prepared in accordance with the procedures  
described in U.S. Patent No. 4,965,392. Then 25  
micrograms of S-benzoylmercaptoacetylglycyl-glycylglycine  
is dissolved in 0.10 ml of 1.0 M carbonate buffer (pH  
12). Then 75 mCi of Tc-99m pertechnetate is added in  
about 1.0 ml followed by 1.0 mg of freshly dissolved  
sodium dithionite (10 mg/ml). This mixture is heated at  
100 °C, plus or minus 4 °C, for 3 minutes, then is cooled  
in an ice bath for 5 minutes to give Tc-99m-MAG<sub>3</sub> as  
determined by ITLC (CH<sub>3</sub>CN solvent); anion exchange HPLC  
(Beckman AX, 10 micron 0.01M Na<sub>2</sub>SO<sub>4</sub>/0.01M Na<sub>3</sub>PO<sub>4</sub>, pH  
7.0); and reverse phase HPLC (Beckman ODS, 5 micron 2%  
CH<sub>3</sub>CN/0.01M Na<sub>3</sub>PO<sub>4</sub>, pH 7.0).

The Tc-99m-MAG<sub>3</sub> complex in carboxylate form is then  
esterified; 0.20 ml 1N HCl, 0.30 ml of 0.2M phosphate  
buffer pH 6.0, 10.0 mg 2,3,5,6-tetrafluorophenol (TFP) in  
0.01 ml 90% CH<sub>3</sub>CN, and 12.5 mg of EDC (1-(3-  
dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride)  
in 0.10 ml of 90% CH<sub>3</sub>CN are combined, and the reactants  
are mixed at room temperature (25 °C, plus or minus 2 °C)  
for 1 hour. At this point, a yield of Tc-99m-MAG<sub>3</sub>-TFP  
ester as determined by ITLC (CH<sub>3</sub>CN solvent); anion  
exchange HPLC (Beckman AX, 10 micron 0.01M Na<sub>2</sub>SO<sub>4</sub>/0.01M  
Na<sub>3</sub>PO<sub>4</sub>, pH 7.0); and reverse phase HPLC (Beckman ODS, 5  
micron 34% CH<sub>3</sub>CN/0.01M Na<sub>3</sub>PO<sub>4</sub>, pH 7.0). The preparation

is purified using a C-18 Baker column. The reaction mixture is, optionally and preferably diluted with 2 ml of water and loaded in the column, washed two times with water and then eight times with 10% C<sub>2</sub>H<sub>5</sub>OH/0.01M Na<sub>3</sub>PO<sub>4</sub>, pH 7.0. The product is eluted with CH<sub>3</sub>CN, and the solvent is removed prior to conjugation with annexin V.

The conjugation of the active ester to annexin V is carried out by adding annexin V in a phosphate buffer, pH 9.5, to the Tc-99m-MAG<sub>3</sub>-TFP ester. The reaction is carried out for at least 30 minutes, and the desired radiolabeled annexin product is obtained by passage through a PD-10 gel filtration column.

### Example III

#### Thrombus Imaging With a Radiolabeled Annexin

##### A. Animal Preparation - LA Vascular Thrombi.

Fasting 25-30 kg Yorkshire swine were sedated with intramuscular Telazol (5-10 mg/kg) (commercially available from AVECO Co.) and commercially available Atropine (1 mg, Elkins-Sinn, Inc., Cherry Hill, NJ). Surital (200 mg) (commercially available from Abbott Laboratories) anaesthesia was administered intravenously. The animals were intubated and given inhalation anaesthesia of 1.5-2% halothane (commercially available from Abbott Laboratories, Abbott Park, IL) and O<sub>2</sub> sufficient to obtain a deep level of anesthesia and physiologic arterial blood gases. Continuous

electrocardiographic monitoring was instituted using alligator clip electrodes. A cut down in the neck region was conducted, and an 8 french catheter (USCI Co, Billerica, MA) was placed in the right common carotid artery for blood pressure and arterial blood gas monitoring as well as for blood sampling.

The swine were placed in a right lateral decubitus position, and a lateral thoracotomy was performed to expose the heart. The incision was held open by a thoracotomy retractor. The pericardium was opened, and the left atrial appendage was isolated from the left atrium by a vascular cross-clamp. Rubber tipped forceps were used to gently crush the appendage. Five minutes later, ricinoleate (1 mg, ICN Pharmaceuticals, Costa Mesa, CA) and thrombin (50 mg, Johnson and Johnson Co., Arlington, TX) were injected into the LAA (left atrial appendage) using a 27 Ga needle. The cross-clamp was removed 10 minutes later.

One hour after the crush injury, Tc-99m-annexin V, prepared in accordance with Example I above, was administered as an intravenous bolus dose in an ear vein. The intravenous line was then flushed with saline. In 7 animals, I-125 labeled ovalbumin was administered as a non-specific control preparable, for example, by the procedure described by Fraker et al., Biochem. Biophys. Res. Commun. 80: 849-57, 1978). Briefly, I-125-radiolabeled ovalbumin was prepared by the Iodogen method, employing 600  $\mu$ g ovalbumin (Sigma Chemical Co.,

St. Louis, MO) and NaI-125 (2mCi, 0.92 nmol).

Image acquisition was performed as described below. At the end of the experimental procedure (generally about 150 minutes), the animals were sacrificed by an intravenous bolus dose of 80 mEq of KCl while the animals were still under general anaesthetic. A final blood sample was taken for well counting. The heart was rapidly excised, washed free of blood and dissected into samples for well counting. Additional samples of carotid artery, lung, liver, spleen, muscle and kidney were obtained in some animals.

B. Controls. Five different types of controls were used: open chest sham; closed chest sham; ovalbumin; indium platelets; and non-specific Tc-99m-labeled antibody.

1. Open Chest Sham. In three animals, the heart was exposed as above, but the left atrium was not isolated, crushed or injected with ricinoleate/thrombin. Marker images with a cobalt marker were performed as described below, and the Tc-99m-annexin V was injected 30-60 minutes after LAA exposure. Imaging and sample acquisition were identical to that described in A above.

2. Closed Chest Sham. In seven animals, an ear intravenous line was established. No thoracotomy was performed. Sedation and anesthesia were identical to that described in A above. The Tc-99m-annexin V (+/- other control radionuclides, such as I-125 ovalbumin or In-111-platelets) was administered and image acquisition

was performed.

3. Ovalbumin. I-125 ovalbumin was administered in 7 animals as a negative control protein. Ovalbumin has a molecular size similar to that of annexin, and exhibits a slightly slower blood clearance.

4. Indium Platelets. In-111 platelet labeling was performed in 7 animals as a positive well counting label. In-111 radiolabeled platelets were prepared in accordance with the procedure described by Stratton et al., Am. J. Cardiol. 47: 874, 1981 and Stratton et al., Circulation 69: 561, 1984. Imaging was not attempted because of the long serum half-life of the In-111-platelets.

5. Non-specific Tc-Labeled Antibody. In a single experiment, a left atrial (LA) thrombi was created by the above method, but Tc-99m-Annexin V was not administered. Instead, a Tc-99m-Fab fragment of an antibody designated as NR-LU-10 was administered. NR-LU-10 is a 150 kilodalton molecular weight IgG2b monoclonal antibody that recognizes an approximately 40 kilodalton glycoprotein antigen expressed on most carcinomas. NR-LU-10 is a well characterized pancarcinoma antibody that has been safely administered to over 565 patients in human clinical trials. NR-LU-10-Fab is prepared in accordance with known techniques and is radiolabeled in accordance with the procedures described in J. Nucl. Med. 32: 1445-51, 1991 and by the modified C-18 Baker purified Tc-99m-N<sub>2</sub>S<sub>2</sub>-TFP procedure described in Example I for preparing radiolabeled annexin V. This Tc-99m-Fab

conjugate was designed as a negative control for both well counting and imaging.

5        C. Imaging. A cobalt marker was placed on the exposed surface of the LAA and held in place with surgical tape affixed to the thoracotomy retractor. The tape was adjusted so that the marker generally moved with the LAA with each cardiac cycle. Marker images were acquired for 10 seconds in each planar view and 10 seconds in each tomographic slice. The cobalt marker was then removed.

10  
15  
20  
25  
A General Electric Starport camera with a general purpose collimator was used to acquire the Tc-99m images. Five minute planar acquisitions were performed sequentially in the left lateral, 45 LAO, and anterior views. These were followed by a 10 minute tomographic acquisition. This full set of 3 planar and 1 tomographic acquisitions was repeated for a total of 5 sets. Care was taken not to move the pig or imaging gantry during the entire imaging sequence. Images were recorded on a Microdelta computer slaved to a VAX mainframe system. Images were stored on tape or on the VAX hard drive. Planar image analysis consisted of first viewing the image with the marker and recording the marker position on the viewing terminal screen. The first image acquired after Tc-99m-annexin V injection was used to define the cardiac blood pool. Each subsequent image was viewed and analyzed using the marker and initial blood pool as references. Each image was scored as positive, equivocal

or negative.

Thirteen animals had left atrial thrombi created and were imaged as described above. Twelve animals had Tc-99m-annexin V injected for imaging, and one animal had a non-specific Tc-99m Fab injected as a control. Closed chest imaging was performed in 7 animals without atrial injury, and open chest sham experiments were performed in 3 animals. In animals with atrial thrombi, all planar images taken at less than 35 minutes after administration of Tc-99m-annexin V were negative. Nine of the atrial thrombi planar images taken at greater than 70 minutes were positive, 1 was equivocal and 2 were negative. All of the tomographic images of atrial thrombi were either positive (n=10) or equivocal (n=2) at imaging times of greater than 2 hours post injection. The average time in which the planar images became positive following administration of Tc-99m-annexin V was 82 minutes (35-135 minutes).

None of the closed chest control animals had positive images. One of the three open chest shams had a positive image at 85 minutes. This false positive is believed to result from the production of surgically-induced thrombi.

These results indicate that intravenous administration of Tc-99m-annexin V permitted acquisition of diagnostic images identifying atrial vascular thrombi within a short time period following conjugate administration.

D. Sample Collection. Samples (both blood and

tissue), as described above, were weighed and placed in vials for immediate counting of Tc-99m. After the Tc-99m had decayed (typically at 5-7 days), the samples were re-assayed to obtain the I-125 counts. Each vial was counted for 1 minute. The counts were corrected for decay, then for weight, and recorded as counts per minute per gram. The counts per minute per gram for each sample were then normalized by dividing the counts per minute per gram of the final blood specimen. Consequently, the results for each sample were calculated in a ratio with the last blood sample, permitting meaningful comparison between animals. This procedure was performed for all radionuclides in a given experiment.

Well counting ratios were obtained for a variety of tissue samples. For the injured tissues and thrombi, multiple specimens were usually taken from the same animal. In those cases, the maximum ratio for any one specimen is reported, as well as the average of all specimens taken. The maximum for each animal was averaged across all animals and is reported as the Maximum Anx-V Ratio.

These results indicate that Tc-99m-annexin V localizes preferentially to atrial thrombi and injured left atrium, with the highest non-target localization occurring in the kidney. The kidney level is at least partially indicative of excretion of Tc-99m-annexin V via the renal route.

Example IV**Radiolabeled Annexin-Galactose Conjugates**

A. Annexin V -Preparation from human placenta and by expression in *E. coli*. Annexin V was purified from human placenta to a final purity of 99% as described by Funakoshi et. al., Biochemistry 26, 5572-78, 1987 and Tait et. al., Biochemistry 27, 6268-76, 1988.

Alternatively, annexin V was obtained, but not employed in the following experiments, by expression in *E. coli*. This expression was conducted as follows.

1. Preparation of bulk fermentation supernatant. Standard molecular biology techniques were used to express annexin V in *E. coli*. More specifically, the protein coding region of annexin V cDNA (see, Funakoshi et. al. referenced above) was inserted into the NdeI/BamHI site of plasmid pET-12a (Novagen, Madison, WI). The selected plasmid was used to transform *E. coli* strain BL21(DE3) (Novagen).

Individual colonies (clones) were isolated on Luria Broth plates containing 50 g/ml ampicillin or carbenicillin, as discussed in Sambrook et al., "Molecular Cloning Laboratory Manual," 2nd ed., Cold Springs Harbor Laboratory Press, 1989. For production, cultures of the selected clones containing the desired plasmid were grown overnight at 37 °C with shaking in Terrific Broth (see, Tartof et. al., Bethesda Res. Lab. Focus, 2: 12, 1987) containing 50 µg/mL carbenicillin (commercially available from Sigma Chemical Co., St.

Louis, Missouri). The culture was diluted 1:20 in the same medium and incubated at 37 °C with shaking for 18-24 hr. The bacteria were harvested by centrifugation, washed once with an equal volume of 0.05 M Tris-HCl, 0.15 M sodium chloride, pH 8, and stored as pellets at -20 °C. Pellets were thawed and resuspended with 10-15% (w/v) cold 0.05 M Tris-HCl, 0.02M Na<sub>4</sub>EDTA, pH 7.2. The suspension was sonicated for 4 min on ice, centrifuged, and the supernatant stored at -70 °C.

2. Purification of annexin V from bulk fermentation supernatant. After thawing, the supernatant was filtered using a 0.1 µm hollow fiber unit (commercially available from A/G Technology, Needham, MA), and the conductivity adjusted with concentrated sodium chloride to equal 0.02 M Tris-HCl, 0.5 M sodium chloride, pH 8.

Polyethyleneimine-WP (J.T. Baker, Phillipsburg, NJ) was suspended in 0.02 M Tris-HCl, 0.5 M sodium chloride, pH 8, and 1 g of resin per 10 ml of bulk fermentation supernatant was added to the filtered supernatant. After stirring for 30 min, the supernatant was removed, exchanged into 0.02 M Tris-HCl, pH 8, and applied to a Q-SEPHAROSE® HP column (Pharmacia, Piscataway, NJ) equilibrated in the same buffer. The column was developed with a step gradient of 0-0.5 M sodium chloride. Fractions were assayed for annexin V by size exclusion HPLC, and fractions containing A<sub>280</sub> absorbing material of appropriate size were pooled. The semi-purified annexin V was exchanged into PBS and

concentrated by ultra-filtration. Gel filtration over SEPHACRYL® 100 HP (Pharmacia) equilibrated in PBS yielded the purified protein. The protein concentration was adjusted to 1 mg/ml, sterile 0.2  $\mu$ m filtered, and stored at 2-8 °C.

B. Derivatization of Annexin V with Galactose. The general method of Lee et al., Biochemistry 15: 6268-76, 1976 was followed.

a. Preparation of 2-imino-2-methoxyethyl-1-thio- -D-galactopyranoside.

A 1.0 M solution of cyanomethyl-2,3,4,6-tetra-O-acetyl-1-thio- -D-galactopyranoside (commercially available from Sigma Chemical Company, St. Louis, MO) was prepared by dissolving 1.83 g (4.5 mmole) in 4.5 ml anhydrous methanol with heating to 50 °C. The solution was maintained at 50 °C, and 0.1 ml (0.45 mmole) of 25% sodium methoxide in methanol (commercially available from Aldrich Chemical Company, Milwaukee, WI) was added with continuous stirring. After 6 hr at 50 °C, the methanol was removed under reduced pressure yielding the galactose methyl imidate as a colorless, viscous oil.

b. Galactosylation of annexin V

The galactose methyl imidate was offered to annexin V in molar ratios of 300:1, 150:1, and 75:1. For the 300:1 offering, 3.35 mg of galactose methyl imidate in methanol was taken to an oil under a stream of nitrogen. To the oil was added 2.53 mg of annexin V in 0.05 M HEPES buffer (commercially available from Sigma Chemical Co., St.

Louis, Missouri), pH 7.4 and 0.04 ml of 0.5 M sodium borate buffer, pH 8.5. The solution was mixed for 1 hr at room temperature by tumbling the reaction vessel, and was then allowed to stand at room temperature overnight (approximately 18 hr). The reaction mixture was diluted with 0.4 ml PBS, transferred to dialysis tubing (6000-8000 MW cut-off), and dialyzed 24 hr against PBS. After removing the material from the dialysis bag, the solution was filtered through a 0.2  $\mu$ m syringe filter. The other offering levels were conducted analogously.

#### C. Characterization of Galactose-Derivatized Annexin

V. Protein concentration was determined using  $A_{280}$  of 0.6 for a 1 mg/ml solution of annexin V. The number of galactose residues per molecule of annexin V was determined by measuring the total number of reactive amines on annexin V before and after reaction with galactose methyl imidate using trinitrobenzenesulfonic acid, as described by Habeeb, Analytical Biochemistry 14: 328-36, 1966.

Offering Ratio	Substitution Ratio
300:1	4.7:1
150:1	2.3:1
75:1	1.1:1

The ability of galactose-modified annexin V to bind to activated platelets was assessed by determining its ability to inhibit the binding of unmodified,  $^{125}$ I radiolabeled annexin V to freshly isolated human

platelets, following the method of Thiagarajan and Tait  
J. Biol. Chem., 265: 17240-43, 1990. The following table  
 shows the results of the competition assay, both in  
 absolute value (left column) and in a value normalized to  
 100% for unmodified annexin V (right column).

Substitution Ratio	Competition (% of control)	Competition (% of control)
4.7	0	0
2.3	5.7	7.6
1.1	21.4	28.7
unmodified	74.5	100

#### D. Preparation of Tc-99m-Annexin V-Galactose.

Annexin V-galactose was radiolabeled with technetium-99m  
 using a diamide dimercaptide ( $N_2S_2$ ) chelating compounds  
 as described by Kasina et. al., J. Nucl. Med. 32: 1445-  
 51, 1991. The preformed active ester chelating compound  
 was diluted with 2.0 ml water and purified before  
 conjugation to annexin V-galactose using a modified  
 conditioned C-18 column (commercially available from J.T.  
 Baker), as described by Fritzberg et. al., Proc. Natl.  
 Acad. Sci. USA 85: 4025-29, 1988, washing with 2.0 ml  
 water eight times followed by drying the column for 5 min  
 and eluting with 100% acetonitrile. The solvent was  
 removed under a steady stream of  $N_2$ . Then 0.15 ml of  
 PBS, 0.35 ml of annexin V-galactose (4.7:1

galactose:annexin V) at 1.0 mg/ml, and 0.2 ml of 0.2M bicarbonate buffer (pH 10.0) were added for conjugation to Tc-99m-N<sub>2</sub>S<sub>2</sub>-TFP ester. After 20 min at room temperature, the Tc-99m-annexin V-galactose conjugate was purified by passage through a G-25 SEPHADEX® PD-10 column (commercially available from Pharmacia) equilibrated with PBS.

Fractions of 1.0 ml were collected, and fractions containing annexin V were pooled. Protein concentration was determined by UV absorption at 280 nm. The radiochemical yield of the conjugate was 35.3%. The radiochemical purity was 99.8%, as assessed by instant thin layer chromatography on silica gel impregnated glass fiber sheets developed in 12% w/v aqueous trichloroacetic acid. The ITLC sheets were cut into two halves, and each half was counted in a gamma counter or a dose calibrator. The radiolabeled annexin V-galactose conjugate precipitated at the origin, and any non-protein-bound radioactivity migrated with the solvent front in this solvent system. Tc-99m-annexin V-galactose conjugate (300-350 g) solution was diluted and stored in PBS containing bovine serum albumin (commercially available from Sigma Chemical Co.) at a final concentration of 15-20 mg/ml PBS prior to intravenous injection.

E. Blood Clearance of Tc-99m-Annexin V and of Tc-99m-Annexin V-Galactose. An animal model, as previously described in Example III, was employed to evaluate blood clearance of the molecules. For N=20 swine, the blood

clearance (Figure 2) is biphasic (biexponential):

$$\%ID/g \quad a = 0.0389 \quad t_{1/2} \quad a = 9.6 \text{ min}$$

$$\%ID/g \quad b = 0.0300 \quad t_{1/2} \quad b = 46 \text{ min}$$

$a + b = 0.0689$ , therefore 57% ( $0.0389/0.0689$ ) of the injected dose/gram of blood is cleared in the faster a phase and 43% of the ID/g is cleared in the slower b phase.

Blood clearance of Tc-99m-annexin V-galactose in the swine (Figure 2) is also biphasic (biexponential):

$$\%ID/g \quad a = 0.0313 \quad t_{1/2} \quad a = 3.5 \text{ min}$$

$$\%ID/g \quad b = 0.0045 \quad t_{1/2} \quad b = 160 \text{ min}$$

$a + b = 0.0358$ , therefore 87% ( $0.0313/0.0358$ ) of the injected dose/gram of blood is cleared very quickly in the early a phase. This more rapid clearance reduces the background radioactivity in the blood compartment, *i.e.*, the noise, and therefore improves the signal to noise ratio. Consequently, thrombus imaging can be achieved at shorter time points.

#### EXAMPLE V

##### Chelating Compound Preparation

4.4 - Diethoxycarbonylpropyl - 1.3 - dianiline: A stirred solution of 2.065 g (1.25 mole) ethyl-4-amino benzoate, 14.35 mL (0.125 mole) 1,3-di-iodopropane and

10.5 g (0.125 mole) sodium bicarbonate in 500 mL dry DMSO is heated at 100°C for 3 hours under nitrogen. Upon cooling, the mixture is poured into 2 L of ice water with stirring, and the resulting precipitate is collected by filtration. The precipitate is then washed with glacial acetic acid (14 x 75 mL) until all of the starting ethyl-4-aminobenzoate has been removed. After drying *in vacuo*, the product, thus obtained, is used in the next step without further purification.

1,3-Di(2-imino-6-ethoxycarbonylbenzthiazolyl-

3-)propane: Ammonium thiocyanate (16.5 g, 0.217 mole) is added to a magnetically stirred suspension of 4,4-diethoxycarbonylpropyl-1,3-dianiline (10.0 g, 0.27 mole) in 1500 mL glacial acetic acid. A solution of bromine (34.6 g, 0.216 mole) in 100 mL glacial acetic acid is then added dropwise to the suspension with stirring at room temperature. After stirring the reaction mixture overnight at room temperature, the dihydrobromide salt of the crude product is collected by filtration and dried.

The product is isolated by dissolving the crude product in hot water, adjusting to basic pH with the addition of saturated sodium bicarbonate solution, collecting the precipitate by filtration, and drying *in vacuo*.

N,N'-Bis(2-disulfidyl-4-hydroxycarbonylphenyl)-1,3-

propyldiamine: Solid potassium hydroxide (20.0 g, 0.357 mole) is added to a suspension of 1,3-di(2-imino-6-ethoxycarbonylbenzthiazolyl-3-)propane (1.0 g, 0.002 mole) in 40 mL distilled water, and the resulting mixture

is heated at 120°C for 12 hours. Complete dissolution occurs after 1 hour. The reaction mixture is then cooled in an ice bath, and the pH is adjusted to 5.0 with 5.0 N acetic acid. The aqueous solution is then extracted with three 100 mL portions of ethyl acetate. The combined ethyl acetate extracts are dried over anhydrous sodium sulfate, and the drying agent is filtered. Removal of solvent yields the product.

To this point, this synthesis is shown schematically in Figure 3.

N,N'-bis(2-disulfidyl-4-hydroxycarbonylphenyl)-1,3-propyldiamine mono-(methyl-aminocaproate) adduct. To a mixture of N,N'-bis(2-disulfidyl-4-hydroxycarbonylphenyl)-1,3-propyldiamine and 2-3 equivalents of methyl 6-aminohexanoate hydrochloride (prepared as described below in Example VI) in dimethylformamide is added 10 equivalents of triethylamine followed by 0.9-1.0 equivalent of BOP (benzotriazol-1-yloxy-tris-(dimethyl-amino)-phosphonium hexafluorophosphate). The mixture is stirred at 15-30°C for 2-24 hours and then concentrated. The residue is diluted with deionized water, and the pH is adjusted to approximately 2 with 1 N aqueous hydrochloric acid. The mixture is again concentrated. The residue is chromatographed on silica gel. The chromatographic fractions containing product are combined and concentrated to afford the product.

This chelating compound is amenable to use with a suitable trifunctional linker or a pair of bifunctional

linkers to form a radiolabeled annexin-galactose cluster conjugate of the present invention. The use thereof with a pair of bifunctional linkers is addressed in the following example.

5

#### EXAMPLE VI

##### **Radiolabeled Annexin-Galactose Cluster Conjugates Bifunctional Linker Approach**

10

#### **A. Preparation of an Eight Galactose Cluster.**

This procedure is schematically shown in Fig. 5.

15

Methyl 6-bromohexanoate. To a 1 L round bottom flask, charged with 20 g (102.5 mmol) of 6-bromohexanoic acid and 500 mL of methanol, was bubbled hydrogen chloride gas for 2-3 minutes. The mixture was stirred at room temperature for 4 hours and concentrated to afford 21.0 g of the product as a yellow oil (99%): <sup>1</sup>H-NMR (200MHz, d<sub>6</sub>-DMSO); 3.57 (s, 3H), 3.51 (t, 2H), 2.30 (t, 2H), 1.78 (pentet, 2H), and 1.62-1.27 (m, 4H) ppm.

20

Methyl 6-aminohexanoate hydrochloride. To a 1 L round bottom flask, charged with 40.0 g aminocaproic acid, was added 500 mL of methanol. Hydrogen chloride gas was bubbled through the mixture for 5 minutes, and the mixture was stirred at room temperature for 5 hours. The mixture was then concentrated via rotary evaporation and then under full vacuum pump pressure (<0.1 mm Hg) to afford 55 g of the product as a white solid (99%): <sup>1</sup>H-NMR (200 MHz, CD<sub>3</sub>OD); 3.67 (s, 3H), 3.02 (t, 2H), 2.68 (s,

25

3H), 2.48 (t, 2H), and 2.03-1.87 (pentet, 2H) ppm.

5        Methyl 6-(trifluoroacetamido)-hexanoate: To a 1 L round bottom flask, charged with 25.0 g (138 mmol) of methyl 6-aminohexanoate hydrochloride and 500 mL of methylene chloride, was added 24 mL (170 mmol) trifluoroacetic anhydride. The mixture was cooled in an ice bath, and 42 mL (301 mmol) of triethylamine was added over a 25-30 minute period. The mixture was stirred at 0°C to room temperature for 2 hours and then concentrated. The residue was diluted with 150 mL of diethyl ether and 150 mL of petroleum ether, and the resulting solution was washed first with 1 N aqueous HCl (3 x 150 mL) and then with saturated aqueous sodium bicarbonate (3 x 150 mL). The organic phase was dried over magnesium sulfate, filtered and concentrated to give 32.9 g of the product as a pale yellow oil (99%): <sup>1</sup>H-NMR (200MHz, d<sub>6</sub>-DMSO); 9.39 (m, 1H), 3.57 (s, 3H), 3.14 (q, 2H), 2.29 (t, 2H), 1.60-1.38 (m, 4H), and 1.32-1.19 (m, 2H) ppm.

20        N,N'-Bis(6-methoxycarbonylhexyl)amine hydrochloride. To a 500 mL dry round bottom flask, charged with 12.0 g (50.0 mmol) of the secondary amide, methyl 6-(trifluoroacetamido)-hexanoate, and 250 mL of dry tetrahydrofuran, was added 2.2 g (55 mmol, 1.1 equiv) of 25        60% sodium hydride. The mixture was stirred at room temperature for 30 minutes and then 10.25 g (49.0 mmol, 0.98 equiv) of the alkyl bromide, methyl 6-bromohexanoate, was added. The mixture was stirred at

reflux for 3 hours, an additional 5.80 g (27.7 mmol, 0.55 equiv) of methyl 6-bromohexanoate was added, and the mixture was stirred at reflux for 70 hours. The mixture was cooled, diluted with 150 mL of 1 N aqueous HCl and then extracted with ethyl acetate (3 x 100 mL). The organic extracts were combined, dried over magnesium sulfate, filtered and concentrated. The residue was diluted with 200 mL of methanol and then treated with 30 mL of 10 N aqueous sodium hydroxide. The mixture was stirred at room temperature for 18 hours and then concentrated. The residue was diluted with 200 mL of deionized water and acidified to pH 1-2 with 37% concentrated HCl. The solution was washed with diethyl ether (3 x 100 mL). The aqueous phase was concentrated. The residue was diluted with 200 mL of methanol and reconcentrated. The subsequent residue was diluted with 250 mL of methanol, and HCl gas was bubbled through for 2-3 minutes followed by stirring at room temperature for 3 hours. The mixture was concentrated. The residue was diluted with 300 mL of methanol and filtered to remove inorganic salts. The filtrate was treated with 3 g of activated charcoal, filtered through Celite (manufactured by J.T. Baker) and concentrated. The residue, an off-white solid, was recrystallized from 100 mL of 2-propanol to afford 7.0 g of the product as a white solid. Concentration of the filtrate and further recrystallization of the residue yielded an additional 1.65 g of the product for a total of 8.65 g (56%):  $^1\text{H}$ -

NMR (200MHz,  $d_6$ -DMSO); 3.57 (s, 3H), 2.90-2.73 (m, 4H), 2.30 (t, 4H), 1.67-1.44 (m, 8H), and 1.37-1.20 (m, 4H) ppm.

5      Methyl 4-methylaminobutyrate hydrochloride. To a 1 L round bottom flask, charged with 30.0 g (195 mmol) of 4-methylaminobutyric acid and 500 mL of methanol, was bubbled HCl gas for 1-2 minutes. The mixture was stirred at room temperature for 3-4 hours and then concentrated to afford 32.5 g of the product as a foamy, off-white solid (99%):  $^1\text{H}$ -NMR (200 MHz,  $\text{CD}_3\text{OD}$ ); 3.67 (s, 3H), 3.03 (t, 2H), 2.68 (s, 3H), 2.48 (t, 2H), and 2.03-1.87 (pentet, 2H) ppm.

10  
15  
20  
25  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2

exchange resin (hydroxide form) commercially available from BioRad, Richmond, CA), and by stirring the solution for 2 hours. The resin was filtered off and washed with 150 mL of deionized water. The aqueous filtrates were combined and concentrated. The residue was diluted with 200 mL of 2-propanol and filtered. The collected solids were rinsed with 100 mL of 2-propanol. The organic filtrates were combined and concentrated. The residue was distilled under reduced pressure to afford 12.85 g of the product as a colorless oil (bp 68°C at 0.1-0.2 mm HG; 64%): <sup>1</sup>H-NMR (200 MHz, D<sub>2</sub>O); 3.52 (t, 2H), 2.56 (t, 2H), 2.31 (s, 3H), and 1.65-1.43 (m, 4H) ppm.

4-(N-Methyl-trifluoroacetamido)-1-butanol. To a 250 mL round bottom flask, charged with 10.0 g (96.9 mmol) of the amine, 4-methylaminobutanol, in 100 mL of dry methanol, was added 17.5 mL (147 mmol) of ethyl trifluoroacetate. The mixture was stirred at room temperature for 24 hours and then concentrated to afford 18.55 g of the product as a near colorless oil (96%): <sup>1</sup>H-NMR (200 MHz, D<sub>2</sub>O); 3.63 and 3.50 (2t's, 4H), 3.20 and 3.05 (d and s, 3H), and 1.82-1.47 (m, 4H) ppm.

1-(p-Toluenesulfonyloxy)-4-(N-methyl-trifluoroacetamido)butane. To a 1 L dry round bottom flask, charged with 17.0 g (85.4 mmol) of the alcohol, 4-(N-methyl-trifluoroacetamido)-1-butanol, in 400 mL of methylene chloride, was added 17.1 g (89.7 mmol, 1.05 equiv) of toluenesulfonyl chloride followed by 30 mL (213 mmol, 2.5 equiv) of triethylamine at 0°C over a 10 minute

period. The mixture was stirred at 0°C to room temperature for 15 hours and then washed with 5% v/v aqueous HCl (3 x 200 mL). The organic phase was dried over magnesium sulfate, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 50:50 hexane/methylene chloride and then with methylene chloride, to give 25.1 g of the product as a pale yellow oil (83%): <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>); 7.80 (d, 2H), 7.37 (d, 2H), 4.07 (m, 2H), 3.41 (m, 3H), 3.09 and 2.98 (q and s, 3H), 2.45 (s, 3H), and 1.68 (m, 4H) ppm: TLC (methylene chloride) R<sub>f</sub> = 0.31.

1-S-(2,3,4,6-tetra-O-acetyl-beta-D-galactopyranosyl)-2-thiopseudourea hydrobromide. To a 250 mL round bottom flask, charged with 5.08 g (60.8 mmol, 1.09 equiv) of thiourea and 36 mL of acetone, was added 25.0 g (66.7 mmol) of tetra-acetyl-alpha-D-galactopyranosyl bromide. The mixture was stirred at reflux for 15-20 minutes and then cooled on ice. The mixture was filtered into a Buchner funnel and rinsed with 25 mL of ice cold acetone. The solids were treated with 50 mL of acetone, refluxed for 15 minutes, cooled on ice, and filtered. The solids were rinsed with 25 mL of cold acetone, air dried and then dried under vacuum to give 22.6 g of the product as a white solid (76%): <sup>1</sup>H-NMR (200MHz, d<sub>6</sub>-DMSO); 9.4-9.0 (broad d, 4H), 5.63 (d, 1H), 5.38 (d, 1H), 5.23 (dd, 1H), 5.09 (t, 1H), 4.40 (t, 1H), 4.04 (dd, 1H), 2.13 (s, 3H), 2.08 (s, 3H), 2.00 (s, 3H), 1.93 (s, 3H) ppm.

4-(N-Methylaminobutyl)-1-thio-beta-D-galactopyranoside. To a 500 mL round bottom flask, charged with 20.7 g (42.5 mmol, 1.07 equiv) of the thiopseudourea hydrobromide prepared as described above in 70 mL of deionized water, was added 6.4 g (46.3 mmol, 1.16 equiv) of potassium carbonate and 4.7 g (45.2 mmol, 1.13 equiv) of sodium bisulfite followed immediately by 14.1 g (39.9 mmol, 1.0 equiv) of the tosylate, 1-(p-toluenesulfonyloxy)-4-(N-methyl-trifluoroacetamido)butane, in 70 mL of acetone. The mixture was stirred at room temperature for 16 hours. The mixture was diluted with 50 mL of brine and extracted with ethyl acetate (3 x 200 mL). The organic extracts were combined, dried over magnesium sulfate, filtered and concentrated. The residue was chromatographed on silica gel, eluting first with 75% methylene chloride/hexane, followed by methylene chloride, then with 2% methanol/methylene chloride and finally with 10% methanol/methylene chloride. Fractions containing alkylation product with different degrees of acetylation were combined and concentrated. The residue was diluted with 250 mL of methanol and 150 mL of deionized water and treated with 110 g of AG-1 X-8 resin (hydroxide form; 2.6 m equiv/g dry weight) commercially available from BioRad. The mixture was stirred at room temperature for 18 hours. The mixture was filtered, and the resin was rinsed with methanol (2 x 150 mL). The filtrates were combined and concentrated to afford 6.1 g of product (54%): <sup>1</sup>H-NMR

(200 MHz, D<sub>2</sub>O); 4.38 (d, 1H), 3.88 (d, 1H), 3.69-3.41 (m, 5H), 2.82-2.64 (m, 4H), 2.43 (s, 3H), and 1.68-1.57 (, 4H) ppm.

5        N-BOC-Bis-methylester. To 1.00 g (3.23 mmol) of the  
amine hydrochloride, N,N-bis-(6-methoxycarbonyl-  
hexyl)amine hydrochloride prepared as described above,  
was added 1.5 mL (10.6 mmol) of triethylamine followed by  
875 mg (3.55 mmol, 1.1 equiv) of BOC-ON, 2-(tert-  
butoxycarbonyloxyimino)-2-phenylacetonitrile. The  
10 mixture was stirred at room temperature for 18 hours and  
then concentrated. The residue was diluted with 100 mL  
of ethyl acetate and washed with 1 N aqueous hydrochloric  
acid (3 x 50 mL), followed by saturated aqueous sodium  
15 bicarbonate (2 x 50 mL). The organic phase was dried  
over magnesium sulfate, filtered and concentrated. the  
residue was chromatographed on silica gel, eluting with  
15% (percentage by volume) ethyl acetate/hexane.  
Chromatographic fractions containing product were  
combined and concentrated to afford 990 mg of product as  
20 a near colorless oil (83%).

25        N-BOC-Bis-acid. To 980 mg (2.62 mmol) of the diester  
prepared in the previous step in 10 mL of methanol was  
added 5.8 mL of 1 N aqueous sodium hydroxide (5.8 mmol).  
The mixture was stirred at room temperature for 16 hours  
and then concentrated. The residue was diluted with 30  
mL of deionized water and acidified to pH 1.5-2. The  
mixture was extracted with ethyl acetate (6 x 50 mL).  
The organic extracts were dried over magnesium sulfate,

filtered and concentrated. The residue was chromatographed on reverse phase C-18 silica gel commercially available from J.T. Baker, eluting with 65% methanol/water. Chromatographic fractions containing product were combined and concentrated to afford 851 mg of product as a near colorless oil (94%).

N-BOC-Tetra-methyl ester. To 825 mg (2.39 mmol) of the bis-acid prepared as described above in 35 mL of dry dimethylformamide, was added 1.75 g (5.65 mmol, 2.36 equiv) of amine hydrochloride, N,N-bis-(6-methoxycarbonylhexyl)amine hydrochloride, and 3.0 mL of triethylamine followed by 2.4 g (5.4 mmol, 2.3 equiv) of BOP. The mixture was stirred at room temperature for 17 hours and then concentrated. The residue was diluted with 100 mL of ethyl acetate and washed with 1 N hydrochloric acid (3 x 50 mL) followed by washing with aqueous sodium bicarbonate (2 x 50 mL). The organic phase was dried over magnesium sulfate, filtered and concentrated. The residue was chromatographed on silica gel, eluting with ethyl acetate. Chromatographic fractions containing product were combined and concentrated to afford 1.63 g of the product as a near colorless oil (80%).

N-BOC-Tetra-acid. To a solution of 1.41 g (1.65 mmol) of tetra-methyl ester prepared as described above in 25 mL of methanol was added 7.4 mL (7.4 mmol) of 1 N aqueous sodium hydroxide. The mixture was stirred at room temperature for 22 hours and then concentrated. The

residue was diluted with 30 mL of deionized water and acidified to pH 2 with 1 N aqueous hydrochloric acid. The mixture was extracted with 3:1 (ratio by volume) ethyl acetate/isopropanol (3 x 100 mL). The organic extracts were concentrated. The residue was chromatographed on reverse phase C-18 silica gel, eluting initially with 50:50 (ratio by volume) methanol/water and eventually with 75:25 methanol/water. Chromatographic fractions containing product were combined and concentrated to afford 1.19 g of the product as a colorless oil (90%).

N-BOC Octa-methyl ester. To a mixture of 501 mg (0.626 mmol) of tetra-acid prepared as described above and 30 mL of dry dimethylformamide was added 968 mg (3.12 mmol, 5.0 equiv) of amine hydrochloride, N,N'-bis-(6-methoxycarboxyhexyl)amine hydrochloride, and 2.0 mL (14.2 mmol) of triethylamine, followed by 1.22 g (2.76 mmol, 4.6 equiv) BOP. The mixture was stirred at room temperature for 19 hours and then concentrated. The residue was diluted with 75 mL of ethyl acetate and washed with 1 N aqueous hydrochloric acid (2 x 50 mL). The organic phase was dried over magnesium sulfate, filtered and concentrated. The residue was chromatographed on reverse phase C-18 silica gel, eluting initially with 60:40 methanol/water and eventually with 90:10 methanol/water. The chromatographic fractions containing product were combined and concentrated to afford 715 mg of the product as a colorless oil (63%).

N-BOC Octa-acid. To a solution of 715 mg (0.393 mmol) of octa-methyl ester prepared as described above in 20 mL of methanol was added 6 mL of 1 N aqueous sodium hydroxide (6 mmol) and 5 mL of deionized water. The mixture was stirred at room temperature for 16 hours and then concentrated. The residue was diluted with 20 mL of deionized water, and the solution was acidified to pH 1.5-2.0. The mixture was concentrated, and the residue was chromatographed on reverse phase C-18 silica gel, eluting initially with 50:50 methanol/water and eventually with 80:20 methanol/water. The chromatographic fractions containing product were combined and concentrated to afford 647 mg of the product as a near colorless oil (96%).

The above procedure is designed for the formation of a galactose cluster of 8 galactose residues. The four galactose version can be made in accordance with this procedure by proceeding from the tetra acid to the galactose derivatization step, which is described below for the 8-galactose cluster. Similarly, 16, 32, etc. galactose cluster constructs can be prepared in accordance with the present invention by introduction of more iterations of the methyl ester and acid formation steps. More specifically, the 16-methyl ester construct, the 16-acid, the 32-methyl ester and so on would be prepared essentially as described above for the tetra and octa forms. When the desired number of acid residues are formed, the galactose derivatization step is employed,

with the proportions of the components adjusted to accommodate the number of acid residues.

N-BOC-Octa-galactosyl construct. To a mixture of 161 mg (94 mmol) of octa-acid prepared as described above and 225 mg (906 micromol, 9.64 equiv) of galactose amine, 4-N-methylaminobutyl)-1-thio-beta-D-galactopyranoside, in 8 mL of dry dimethylformamide was added 0.5 mL (3.54 mmol) of triethylamine followed by 371 mg (839 micromol, 8.4 equiv) of BOP. The mixture was stirred at room temperature for 17 hours and then concentrated. The residue was chromatographed on reverse phase C-18 silica gel, eluting initially with 40:60 methanol/water and finally with 70:30 methanol/water. The chromatographic fractions containing product were combined and concentrated to afford 170 mg of the product as a near colorless oil (47%).

Octa-galactosyl amine. To 170 mg of the N-BOC-octa-galactosyl construct prepared as described above was added 5 mL of trifluoroacetic acid. The mixture was stirred at room temperature for 10 minutes and then concentrated. The residue was diluted with 10 mL of methanol and reconcentrated. The residue is used without further purification.

#### B. Extender-Galactose Cluster Preparation.

This procedure is schematically shown in Fig. 6.

Methyl 6-(N-BOC)-aminocaproate. To a mixture of amine hydrochloride, methyl-6-aminohexanoate hydrochloride, prepared as described above is added 1.1

equivalents of BOC-ON followed by 2-3 equivalents of triethylamine. The mixture is stirred at 15-30°C for 16-24 hours and the concentrated. The residue is dissolved in ethyl acetate and washed with 1 N aqueous hydrochloric acid and then with saturated aqueous sodium bicarbonate. The organic phase is dried over magnesium sulfate, filtered and concentrated via reduced pressure rotary evaporation. The residue is chromatographed on silica gel, eluting with 25% ethyl acetate/hexane. The chromatographic fractions containing the product are combined and concentrated to afford the product.

6-(N-BOC)-aminocaproic acid. To a solution of the methyl ester, methyl 6-(N-BOC)-aminocaproate, in methanol is added 1.5 equivalents of 1 N aqueous sodium hydroxide. The mixture is stirred at 15-30°C for 16-24 hours and then concentrated. The residue is diluted with deionized water and extracted with ethyl acetate. The organic extracts are combined, dried over magnesium sulfate, filtered and concentrated. The residue is chromatographed on silica gel, eluting initially with 25% ethyl acetate/hexane and finally with 100% ethyl acetate. The chromatographic fractions containing the product are combined and concentrated to afford the product.

N-BOC extended octa-galactosyl construct. To a solution of the octa-galactosyl amine prepared as described above in dimethylformamide and 1.5-3 equivalents of 6-(N-BOC)-aminocaproic acid is added 4-6 equivalents of triethylamine followed by 1.1-1.5

equivalents of BOP. The mixture is stirred at 15-30°C for 4-24 hours and then concentrated. The residue is diluted with deionized water, and the pH is adjusted to 1.5-2.0 by addition of 1 N aqueous hydrochloric acid. The mixture is washed with ethyl acetate. The aqueous phase is concentrated, and the residue is chromatographed on reverse phase C-18 silica gel, eluting initially with 50:50 methanol/water and finally with 65:35 methanol/water. The chromatographic fractions containing product are combined and concentrated to afford the product.

Amine extended octa-galactosyl construct. To the N-BOC protected amine prepared in the previous step is added trifluoroacetic acid. The mixture is stirred at 15-30°C for 10 minutes and then concentrated. The residue is diluted with methanol and reconcentrated to afford the product which is used without further purification.

#### C. Conjugation of Galactose Cluster with Chelating Compound and Annexin Components.

Octa-galactosyl-chelating compound construct. To a mixture of amine extended octa-galactosyl amine, 1.2 equivalents of N,N'-bis(2-disulfidyl-4-hydroxycarbonylphenyl)-1,3-propyldiamine mono-(methyl-aminocaproate) adduct, and 5 equivalents of triethylamine in dimethylformamide is added 1.1 equivalents of BOP. The mixture is stirred at 15-30°C for 2 to 24 hours and then concentrated. The residue is diluted with deionized

water, and the pH is adjusted to 2.5 by the addition of 1 N aqueous hydrochloric acid. The mixture is washed with ethyl acetate. The aqueous phase is concentrated. The residue is chromatographed on reverse phase C-18 silica gel. The fractions containing the product are combined and concentrated to give the product.

Octa-galactosyl-chelating compound carboxylic acid construct. To a methanolic solution of the ester bearing octa-galactosyl-chelating compound construct prepared previously is added 4-5 equivalents of 1 N aqueous sodium hydroxide. The mixture is stirred at 15-30°C for 16-24 hours. The mixture is concentrated, and the residue is diluted with deionized water. The pH of the resulting solution is adjusted to approximately 2.5, and the solution is reconcentrated. The residue is chromatographed on reverse phase C-18 silica gel. The fractions containing the product are combined and concentrated to afford the product.

Annexin V-chelating compound-octa-galactosyl construct. The octa-galactosyl-chelating compound carboxylic acid construct is offered to annexin V in molar ratios of 30:1, 15:1, 5:1 and 2:1. The conjugation of annexin V is carried out via activation of the carboxylic acid functional group of the galactose cluster-chelating compound construct with benzotriazol-1-yl-oxy-tris-(dimethylamino)phosphonium hexafluorophosphate (BOP). To the galactose cluster-chelating compound construct in DMF, an equimolar

concentration of BOP in DMF is added. Annexin V in 0.05 M HEPES buffer (commercially available from Sigma Chemical Co., St. Louis, MO), pH 7.4 is added to the reaction mixture followed by addition of 0.5 M borate buffer, pH 8.5. The DMF concentration in the reaction mixture is maintained between 5-20%. The solution is mixed for 1 hour at room temperature by tumbling the reaction vessel, and is then allowed to stand at room temperature overnight (approximately 18-24 hours). The reaction mixture is then diluted with PBS, transferred to dialysis tubing (6000-8000 molecular weight cut off), and dialyzed for 24 hours against PBS. After removing the material from the dialysis bag, the solution is filtered through a 0.2 micrometer syringe filter. All of the offering levels are conducted analogously.

#### D. Characterization of Galactose Cluster-Chelating Compound-Annexin V Conjugate.

Protein concentration is determined using  $A_{280}$  of 0.6 for a 1 mg/mL solution of annexin V. The number of galactose residues per molecule of annexin V is determined by measuring the total number of reactive amines on Annexin V before and after reaction with galactosyl cluster-chelating compound conjugate using trinitro-benzenesulfonic acid, as described by Habeeb, Analytical Biochemistry, 14: 328-36, 1966. The ability of galactose-chelating compound-annexin V to bind to activated platelets is assessed by determining its

ability to inhibit the binding of unmodified, I-125-radiolabeled annexin V to freshly isolated human platelets, following the method of Thiagarajan and Tait, J. Biol. Chem., 265: 17,240-43, 1990.

5 E. Radiolabeling Procedure for use in Post-Formed Chelate Conjugation Method.

100  
150  
160  
170  
180  
190  
200  
210  
220  
230  
240  
250  
260  
270  
280  
290  
300  
310  
320  
330  
340  
350  
360  
370  
380  
390  
400  
410  
420  
430  
440  
450  
460  
470  
480  
490  
500  
510  
520  
530  
540  
550  
560  
570  
580  
590  
600  
610  
620  
630  
640  
650  
660  
670  
680  
690  
700  
710  
720  
730  
740  
750  
760  
770  
780  
790  
800  
810  
820  
830  
840  
850  
860  
870  
880  
890  
900  
910  
920  
930  
940  
950  
960  
970  
980  
990  
1000  
1010  
1020  
1030  
1040  
1050  
1060  
1070  
1080  
1090  
1100  
1110  
1120  
1130  
1140  
1150  
1160  
1170  
1180  
1190  
1200  
1210  
1220  
1230  
1240  
1250  
1260  
1270  
1280  
1290  
1300  
1310  
1320  
1330  
1340  
1350  
1360  
1370  
1380  
1390  
1400  
1410  
1420  
1430  
1440  
1450  
1460  
1470  
1480  
1490  
1500  
1510  
1520  
1530  
1540  
1550  
1560  
1570  
1580  
1590  
1600  
1610  
1620  
1630  
1640  
1650  
1660  
1670  
1680  
1690  
1700  
1710  
1720  
1730  
1740  
1750  
1760  
1770  
1780  
1790  
1800  
1810  
1820  
1830  
1840  
1850  
1860  
1870  
1880  
1890  
1900  
1910  
1920  
1930  
1940  
1950  
1960  
1970  
1980  
1990  
2000  
2010  
2020  
2030  
2040  
2050  
2060  
2070  
2080  
2090  
2100  
2110  
2120  
2130  
2140  
2150  
2160  
2170  
2180  
2190  
2200  
2210  
2220  
2230  
2240  
2250  
2260  
2270  
2280  
2290  
2300  
2310  
2320  
2330  
2340  
2350  
2360  
2370  
2380  
2390  
2400  
2410  
2420  
2430  
2440  
2450  
2460  
2470  
2480  
2490  
2500  
2510  
2520  
2530  
2540  
2550  
2560  
2570  
2580  
2590  
2600  
2610  
2620  
2630  
2640  
2650  
2660  
2670  
2680  
2690  
2700  
2710  
2720  
2730  
2740  
2750  
2760  
2770  
2780  
2790  
2800  
2810  
2820  
2830  
2840  
2850  
2860  
2870  
2880  
2890  
2900  
2910  
2920  
2930  
2940  
2950  
2960  
2970  
2980  
2990  
3000  
3010  
3020  
3030  
3040  
3050  
3060  
3070  
3080  
3090  
3100  
3110  
3120  
3130  
3140  
3150  
3160  
3170  
3180  
3190  
3200  
3210  
3220  
3230  
3240  
3250  
3260  
3270  
3280  
3290  
3300  
3310  
3320  
3330  
3340  
3350  
3360  
3370  
3380  
3390  
3400  
3410  
3420  
3430  
3440  
3450  
3460  
3470  
3480  
3490  
3500  
3510  
3520  
3530  
3540  
3550  
3560  
3570  
3580  
3590  
3600  
3610  
3620  
3630  
3640  
3650  
3660  
3670  
3680  
3690  
3700  
3710  
3720  
3730  
3740  
3750  
3760  
3770  
3780  
3790  
3800  
3810  
3820  
3830  
3840  
3850  
3860  
3870  
3880  
3890  
3900  
3910  
3920  
3930  
3940  
3950  
3960  
3970  
3980  
3990  
4000  
4010  
4020  
4030  
4040  
4050  
4060  
4070  
4080  
4090  
4100  
4110  
4120  
4130  
4140  
4150  
4160  
4170  
4180  
4190  
4200  
4210  
4220  
4230  
4240  
4250  
4260  
4270  
4280  
4290  
4300  
4310  
4320  
4330  
4340  
4350  
4360  
4370  
4380  
4390  
4400  
4410  
4420  
4430  
4440  
4450  
4460  
4470  
4480  
4490  
4500  
4510  
4520  
4530  
4540  
4550  
4560  
4570  
4580  
4590  
4600  
4610  
4620  
4630  
4640  
4650  
4660  
4670  
4680  
4690  
4700  
4710  
4720  
4730  
4740  
4750  
4760  
4770  
4780  
4790  
4800  
4810  
4820  
4830  
4840  
4850  
4860  
4870  
4880  
4890  
4900  
4910  
4920  
4930  
4940  
4950  
4960  
4970  
4980  
4990  
5000  
5010  
5020  
5030  
5040  
5050  
5060  
5070  
5080  
5090  
5100  
5110  
5120  
5130  
5140  
5150  
5160  
5170  
5180  
5190  
5200  
5210  
5220  
5230  
5240  
5250  
5260  
5270  
5280  
5290  
5300  
5310  
5320  
5330  
5340  
5350  
5360  
5370  
5380  
5390  
5400  
5410  
5420  
5430  
5440  
5450  
5460  
5470  
5480  
5490  
5500  
5510  
5520  
5530  
5540  
5550  
5560  
5570  
5580  
5590  
5600  
5610  
5620  
5630  
5640  
5650  
5660  
5670  
5680  
5690  
5700  
5710  
5720  
5730  
5740  
5750  
5760  
5770  
5780  
5790  
5800  
5810  
5820  
5830  
5840  
5850  
5860  
5870  
5880  
5890  
5900  
5910  
5920  
5930  
5940  
5950  
5960  
5970  
5980  
5990  
6000  
6010  
6020  
6030  
6040  
6050  
6060  
6070  
6080  
6090  
6100  
6110  
6120  
6130  
6140  
6150  
6160  
6170  
6180  
6190  
6200  
6210  
6220  
6230  
6240  
6250  
6260  
6270  
6280  
6290  
6300  
6310  
6320  
6330  
6340  
6350  
6360  
6370  
6380  
6390  
6400  
6410  
6420  
6430  
6440  
6450  
6460  
6470  
6480  
6490  
6500  
6510  
6520  
6530  
6540  
6550  
6560  
6570  
6580  
6590  
6600  
6610  
6620  
6630  
6640  
6650  
6660  
6670  
6680  
6690  
6700  
6710  
6720  
6730  
6740  
6750  
6760  
6770  
6780  
6790  
6800  
6810  
6820  
6830  
6840  
6850  
6860  
6870  
6880  
6890  
6900  
6910  
6920  
6930  
6940  
6950  
6960  
6970  
6980  
6990  
7000  
7010  
7020  
7030  
7040  
7050  
7060  
7070  
7080  
7090  
7100  
7110  
7120  
7130  
7140  
7150  
7160  
7170  
7180  
7190  
7200  
7210  
7220  
7230  
7240  
7250  
7260  
7270  
7280  
7290  
7300  
7310  
7320  
7330  
7340  
7350  
7360  
7370  
7380  
7390  
7400  
7410  
7420  
7430  
7440  
7450  
7460  
7470  
7480  
7490  
7500  
7510  
7520  
7530  
7540  
7550  
7560  
7570  
7580  
7590  
7600  
7610  
7620  
7630  
7640  
7650  
7660  
7670  
7680  
7690  
7700  
7710  
7720  
7730  
7740  
7750  
7760  
7770  
7780  
7790  
7800  
7810  
7820  
7830  
7840  
7850  
7860  
7870  
7880  
7890  
7900  
7910  
7920  
7930  
7940  
7950  
7960  
7970  
7980  
7990  
8000  
8010  
8020  
8030  
8040  
8050  
8060  
8070  
8080  
8090  
8100  
8110  
8120  
8130  
8140  
8150  
8160  
8170  
8180  
8190  
8200  
8210  
8220  
8230  
8240  
8250  
8260  
8270  
8280  
8290  
8300  
8310  
8320  
8330  
8340  
8350  
8360  
8370  
8380  
8390  
8400  
8410  
8420  
8430  
8440  
8450  
8460  
8470  
8480  
8490  
8500  
8510  
8520  
8530  
8540  
8550  
8560  
8570  
8580  
8590  
8600  
8610  
8620  
8630  
8640  
8650  
8660  
8670  
8680  
8690  
8700  
8710  
8720  
8730  
8740  
8750  
8760  
8770  
8780  
8790  
8800  
8810  
8820  
8830  
8840  
8850  
8860  
8870  
8880  
8890  
8900  
8910  
8920  
8930  
8940  
8950  
8960  
8970  
8980  
8990  
9000  
9010  
9020  
9030  
9040  
9050  
9060  
9070  
9080  
9090  
9100  
9110  
9120  
9130  
9140  
9150  
9160  
9170  
9180  
9190  
9200  
9210  
9220  
9230  
9240  
9250  
9260  
9270  
9280  
9290  
9300  
9310  
9320  
9330  
9340  
9350  
9360  
9370  
9380  
9390  
9400  
9410  
9420  
9430  
9440  
9450  
9460  
9470  
9480  
9490  
9500  
9510  
9520  
9530  
9540  
9550  
9560  
9570  
9580  
9590  
9600  
9610  
9620  
9630  
9640  
9650  
9660  
9670  
9680  
9690  
9700  
9710  
9720  
9730  
9740  
9750  
9760  
9770  
9780  
9790  
9800  
9810  
9820  
9830  
9840  
9850  
9860  
9870  
9880  
9890  
9900  
9910  
9920  
9930  
9940  
9950  
9960  
9970  
9980  
9990  
10000

Method A: Stannous gluconate kits are prepared containing 5 mg sodium gluconate, 100 micrograms stannous chloride, 1.0 mg (1 mg/mL) of galactose cluster-chelating compound-annexin V, and 0.1 to 1.0 mg of lactose. The pH is maintained between 5 and 7 using HCl, acetic acid or NaOH. To the stannous gluconate kit is added 1.0mL sodium pertechnetate (Tc-99m) with a specific activity of about 50 mCi. The vial is incubated at 25-37°C for 5-30 minutes. The percent formation of labeled conjugate, remaining TcO<sub>4</sub>, and hydrolyzed reduced technetium is determined by ITLC in 12% TCA as developing solvent.

20 Method B: Stannous tartrate kits are prepared in an evacuator vial under nitrogen to contain 0.5 mL of disodium tartrate (10 mg/mL) and 0.1 mL stannous chloride (1.0 mg/mL in ethanol). The pH of the solution is kept between 5 and 7, preferably 6.0. To this stannous tartrate solution is added 1.0 mL of sodium pertechnetate (50 mCi), and the solution is allowed to stand at room temperature. In an evacuated vial, 200 microliters of sodium phosphate (0.5 M, pH 8 or 10) and 1.0 mL of galactose cluster-chelating compound-annexin V conjugate (1.0 mg/mL) are added successively. Then Tc-99m-tartrate

25

(50 mCi) is added, and the vial is incubated at 25-37°C for 5-30 minutes. The percent formation of labeled conjugate, remaining  $\text{TcO}_4$ , and hydrolyzed reduced technetium is determined by ITLC in 12% (w/v) trichloroacetic acid as developing solvent.

Constructs prepared in accordance with this Example are tested in accordance with the procedures set forth above (Example III and Example IV(E)) to verify usefulness in clot imaging applications, for example.

#### EXAMPLE VII

##### **Annexin-Galactose Cluster Conjugates Trifunctional Linker Approach**

###### **A. Chelating Compound Preparation.**

Production of chelating compound N,N'-bis(2-disulfidyl-4-methylphenyl)-gamma,gamma'-diamino-isovalerate N-hydroxysuccinimide, as shown schematically in Figure 4.

3-Iodomethyl-4-iodobutyric acid: To a solution of 1.61 g (10 mmole) 3-hydroxymethyl-4-butanolide (prepared by the procedure of Kinoshita and Hirano, J. Heterocyclic Chem., 29: 1025, 1992) in 100 mL carbon tetrachloride is added 8 g (40 mmole) of iodotrimethylsilane. The reaction mixture is heated at 50°C for 12 hours under nitrogen. The mixture is diluted with chloroform and washed with water (3 x 100 mL), 5% aqueous sodium thiosulfate (100mL), 10% aqueous sodium bicarbonate and

brine. The organic layer is dried over magnesium sulfate, filtered and evaporated to give the desired crude product. The crude product is purified by silica gel chromatography (ethyl acetate-hexane = 3:7 as the eluting solvent) to give 3-iodomethyl-4-iodobutyric acid.

Ethyl-3-iodomethyl-4-iodobutyrate: A solution of 2.831 g (8 mmole) 3-iodomethyl-4-iodobutyric acid in 80 mL ethanol is saturated with HCl gas at 0°C. After stirring the solution at room temperature for two days, the solvent is removed under vacuum, and the residue is dissolved in dichloromethane. The dichloromethane layer is washed with 10% aqueous sodium bicarbonate (3 x 100 mL), water (1 x 100 mL) and brine. The separated dichloromethane layer is dried over with magnesium sulfate, filtered and evaporated to give ethyl-3-iodomethyl-4-iodobutyrate.

Ethyl-gamma, gamma'-di(4-methylanilino) isovalerate: A stirred solution of 7.5 g (70 mmole) 4-toluidine, 2.764 g (7 mmole) ethyl-3-iodomethyl-4-iodobutyrate and 0.588 g (7 mmole) sodium bicarbonate in 30 mL dry dimethyl sulfoxide is heated at 100°C for 3 hours under nitrogen. The cooled mixture is poured onto 400 mL ice water with stirring. The resulting precipitate is collected by filtration. The remaining 4-toluidine in the precipitate is removed by washing with aqueous acetic acid several times. The product is obtained by recrystallization of the washed precipitate in heptane.

Ethyl-gamma, gamma'-[1,3-di(2-imino-6-methyl

benzthiazolyl-3)]isovalerate: To a magnetically stirred suspension of 2.0 g (6.5 mmole) ethyl-gamma,gamma'-di(4-methylanilino)isovalerate in 250 mL glacial acetic acid is added ammonium thiocyanate (3.5 g, 0.046 mole) followed by the dropwise addition of a solution of bromine (7.27 g, 0.046 mole) in 50 mL glacial acetic acid. After addition is complete, stirring is continued overnight. The yellow precipitate of dihydrobromide salt is filtered and dried. The dried solid is then dissolved in hot water and the benzothiazole free base is liberated with saturated sodium bicarbonate solution. The white solid is filtered and dried to give crude product which is used without further purification.

N,N'-Bis(2-disulfidyl-4-methylphenyl)-gamma, gamma'-diaminoisovaleric acid: To a suspension of ethyl-gamma,gamma'-[1,3-di(2-imino-6-methyl benzthiazolyl-3)]isovalerate in 40 mL distilled water, solid potassium hydroxide pellets (20.0 g, 0.037 mole) are added, and the resulting solution is heated at 120°C for 15-24 hours. After several hours of heating, the suspension becomes a clear solution. The reaction mixture is cooled in an ice bath and acidified with 5.0 N acetic acid to pH 5.0, and the aqueous solution is extracted with three 100 mL portions of ethyl acetate. The combined ethyl acetate extracts are dried over anhydrous sodium sulfate and filtered. Solvent from the filtrate is removed under reduced pressure to give crude product. This crude product is chromatographed on silica gel column using a

20:80 mixture of ethyl acetate:hexane with 1% acetic acid as eluting solvent to give the product as a crystalline yellow solid.

N,N'-Bis(2-disulfidyl-4-methylphenyl)-gamma, gamma'-diaminoisovalerate N-hydroxysuccinimide: N,N'-Bis(2-disulfidyl-4-methylphenyl)-gamma, gamma'-diaminoisovaleric acid is reacted with N-hydroxysuccinimide (NHS) and dicyclohexylcarbodiimide (DCC) in either tetrahydrofuran (THF) or dimethylformamide (DMF) at room temperature. After stirring overnight at room temperature, the solvent is removed, and the crude product is purified by column chromatography on silica gel.

#### B. Conjugate formation.

This chelating compound is amenable to use with a suitable trifunctional linker to form a radiolabeled annexin- galactose cluster conjugate of the present invention as described below.

Commercially available N-epsilon-t-BOC-lysine (Sigma Chemical Company) is converted, using trifluoroacetic anhydride, to its N-alpha-trifluoroacetamide adduct. Activation of the carboxylic acid functionality, for example with BOP (benzotriazol-1-yloxy-tris(dimethyl-amino)-phosphonium hexafluorophosphate) commercially available from Aldrich Chemical Co., and reaction of the activated moiety with the single available amine on a galactose cluster, e.g., formed as described above, affords a galactose cluster-trifunctional linker species.

The alpha-amine of lysine trifunctional linker component  
 of the galactose cluster-trifunctional linker species is  
 deblocked using methanolic sodium hydroxide. Reaction  
 with the N-hydroxysuccinimide ester of the chelating  
 compound molecule formed as set forth in part A of this  
 example affords a galactose cluster-chelating compound  
 trifunctional linker species. Deprotection of the  
 epsilon amine of the lysine trifunctional linker  
 component using trifluoroacetic acid, followed by  
 reaction with succinic anhydride provides an available  
 carboxylic acid functionality through which the annexin  
 may be conjugated following activation of the carboxylic  
 acid (e.g., with BOP).

#### EXAMPLE VIII

##### Method for Producing a Cell Expression Clone of Annexin V

A parent clone, HPAP1.6, is described in Funakoshi et  
 al., "Primary Structure of Human Placental Anticoagulant  
 Protein", *Biochemistry*, Vol. 26, pp. 8087-8092 (1987).  
 Polymerase chain reaction (PCR) was used to amplify the  
 annexin gene from the  $\lambda$ HPAP1.6 parent clone. The sense  
 primer (CAT ATG GCA CAG CTT CTC A) contained an NdeI  
 restriction site (underlined) and the first 16  
 nucleotides of the annexin leader sequence, beginning  
 with the ATG start codon. The antisense primer (GGA TCC  
 TTA GTC ATC TTC TCC ACA) encoded the end of the coding  
 sequence, a stop codon (bold) and BamHI restriction site

(underlined). The PCR product and plasmid pET-12a (Figure 2) obtained from Novagen (Palo Alto, California) were each digested with NdeI and BamHI and ligated together with T4 DNA ligase. A portion of the ligation solution was transformed into an *E. Coli* host strain and selected on nutrient agar plates containing ampicillin. Plasmid from the resultant clone was designated pET-12a-PAP1-E287G, 7/16/93, clone 1. Dideoxy DNA sequence analysis showed that this plasmid contained DNA that matched the wild-type annexin sequence except for two mutations [855 G → A (silent); 860 A → G (converts Glu-287 to Gly-287)].

The sequence changes present in pET-12a-PAP-E287G, 7/16/93, clone 1 were corrected in the following manner. The plasmid was digested with restriction enzymes SfuI and BamHI which excised a fragment of approximately 240 base pairs containing both mutations. The fragment was replaced with a 240 base pair fragment from an independent clone known to contain the wild-type sequence. The DNAs were ligated and transformed into an *E. Coli* host. *E. Coli* colonies containing plasmid DNA were selected on ampicillin containing nutrient agar plates. The resulting clone harbored plasmid pET-12a-PAP1, 3/7/94, clone 1 (Figure 3). DNA sequencing confirmed that the annexin coding sequence on the plasmid matched the wild-type sequence exactly.

The host strain BL21(DE3), received from Novagen, was transformed with the plasmid pET-12A-PAP1, 3/7/94, clone

1. A glycerol stock was made of the resulting transformant BL21(DE3) (pET-12A-PAP1, 3/7/94, clone 1) and stored at  $\leq -65^{\circ}\text{C}$ .

Growth of *E. coli* BL21(DE3) (pET-12A-PAP1, 3/7/94, clone 1) in liquid culture at  $37^{\circ}\text{C}$  resulted in accumulation of the Annexin V protein in the *E. coli* cytoplasmic space.

#### EXAMPLE IX

#### Procedure for Modification (Replacement of an amino acid) of Annexin V

The plasmid pET-12a is described above in Example VIII. Annexin amino acid variant genes were placed between the NdeI and BamHI restriction sites on the pET-12a plasmid.

An independent modification of annexin V was created by site-directed amino acid alteration, utilizing PCR amplification. To accomplish this alteration, oligonucleotides which anneal in the antisense direction of the 3' end of annexin were created. The particular alteration of annexin V, described herein, is a replacement of a cysteine residue (position 316) with alanine, or another non-sulfur containing amino acid, such as serine. In order to achieve this alteration, the nucleotide sequence was changed from TGT to GCA. The oligonucleotide includes a BamHI restriction enzyme site and a stop codon.

The sense oligonucleotide anneals within the T7 promoter region of pET-12a, just upstream from the NdeI restriction site or the 5' end of annexin V. The following are the anti-sense and sense oligonucleotide sequence, respectively:

(Nx168 Cys TO Ala antisense)

5' GTACCTGGATCCTTAGTCATCTTCTCCGCGAGCAGCAGAAGAGCTTTCTT 3';  
and (T7 sense)

5' CGAAATTAATACGACTCACTATAGGG 3'.

The Nx168 and T7 oligonucleotides were synthesized using DNA synthesizer, model 381A (Applied Biosystem Inc., model 381A, Foster City, CA). After synthesis was complete both oligonucleotides were deprotected as per appendix 5 of the above manufacturer's protocol.

Purification was done using Sephadex™ G-25 column (Pharmacia, Uppsala, Sweden).

PCR reactions were done using ULTma™ polymerase (Perkin Elmer, Norwalk, CT). The reactions were performed using approximately 10ng of pET12a-annexin V template, 30 pMol of T7 and Nx168 oligonucleotides, and 0.4 mM of each nucleotide. Mineral oil overlay was included. The reactions were placed in a Coy temperature cyclor model 110P and incubated for five minutes at 94°C. Approximately 2 minutes into the incubation, 0.5 units of polymerase was added. The reactions were then cycled 30 times at 94°C for 30 seconds, 55°C for one minute, and 74°C for one minute. The reactions were then incubated for 5 minutes at 74°C, and then soaked at 15°C.

The resulting annexin-ala substituted gene was then purified using a DNA purification kit (Promega Magic™ PCR Preps DNA Purification System, Madison, WI) according to manufacturer's protocol.

5           Annexin-ala and pET-12a were then digested with NdeI and BamHI (Gibco/BRL, Gaithersburg, MD) restriction enzymes. Digested products were then purified by agarose gel electrophoresis. Appropriate bands were excised and purified using GeneClean II® Kit (Bio 101, Inc., La Jolla, CA) according to manufacturer's protocol. The annexin-ala and pET-12a were then ligated with T4 DNA ligase (Promega) for 12-18 hours at 4°C.

10  
15  
20           Annexin-ala-pET-12a were transformed into *E. coli* strain DH5α Max. efficiency competent cells (Gibco/BRL). Any other suitable host strain for the pET-12a plasmid may also be used. The *E. coli* transformed cells are placed on LB agar (Gibco/BRL), supplemented with 100 ug/ml ampicillin (International Biotechnologies Inc., New Haven, CT) and incubated for 12-18 hours at 37°C. A suitable colony, was then selected for placement in Terrific Broth (Gibco/BRL), which has been supplemented with 100 ug/ml ampicillin (International Biotechnologies Inc.)

25           Several colonies from the LB agar plates were selected and expanded. The plasmid DNA is purified using Promega's Wizard miniprep purification kit. Once plasmid DNA is purified, colonies are screened by sequence analysis in order to confirm that a particular clone

selected contains the amino acid modification or alteration.

#### EXAMPLE X

##### 5 Preparation and Radiolabeling of a Modified Annexin

The modified annexin, wherein accessible sulfhydryl groups are added to the N-terminus of the annexin molecule, can be prepared in accordance with the procedures set forth in Tanaka, et al., Biochem., 35: 922-929, 1996, as follows.

Preparation of N-Terminally Extended Annexin V  
(Annexin V-N-6, also referred to as modified annexin).  
First, a mutant annexin VcDNA (pANKVC-S) was designed in which the Cys<sub>116</sub> codon was replaced by a Ser codon, an NdeI site was introduced prior to the initiator Met codon, and a BamHI site was introduced after the stop codon. This was constructed by PCR using an annexin V cDNA (pPAP-I-1.6; Funakoshi et al., 1987 Biochemistry 26:5572-5578, Cookson et al., 1994 Genomics 20:463-467) as a template and two oligonucleotides, 5'-  
ggaattc·cat·atg·gca·cag·gtt·ctc·aga·ggc·act·gtg-3' and 5'-  
cgc·gga·tcc·tta·gtc·atc·ttc·tcc·gga·gag·cag-3. This DNA was ligated into the NdeI and BamHI cloning sites in the pET12a plasmid (Novagen, Madison, WI). An oligonucleotide (5'-t·atg·gca·tgt·gac·cat·tc-3') and its inverse complement (5'-t·aga·atg·gtc·aca·tgc·ca-3') were prepared to encode six amino acid residues (Met-Ala-Cys-

Asp-His-Ser) with *Nde*I-compatible overhangs at both ends. The two complementary oligonucleotides were annealed and the product was ligated into pANXVC-S that was digested with *Nde*I to produce plasmid pANXVC-S-N6. DNA sequencing of this plasmid confirmed that the intended mutations had been correctly introduced.

This plasmid (pANXVC-S-N6) was then transformed into *Escherichia coli* strain K-38 containing the pGP-1 plasmid for production of recombinant protein by expression from the T7 promotor of the vector. The cells were grown at 30°C in 3 L of 2% L-broth in 50 mM potassium phosphate, pH 7.4, 25 µM kanamycin. After 4 h ( $OD_{600}=0.4$ ), cells were given a heat shock by placing culture flasks in a 42°C water bath for 20 min and the cultivation was continued at 37°C for 2 h. Cells were then harvested and stored frozen overnight at -20°C.

Thawed cells were suspended in 100 mL of 50 mM Tris-HCl, pH 8.0, 10 mM  $CaCl_2$ , 0.1 mM DFP containing 1 mg of leupeptin and 1 mg of pepstatin and sonicated for 5 min. The cell lysate was treated with 5 µL of DNase (10 units/mL) for 10 min at room temperature. Then, pellets were collected by centrifugation, and annexin V-N6 was extracted from the pellets by stirring for 1 h at 4°C in 70 mL of 50 mM Tris-HCl, pH 8.0, 8 M urea, 0.1 M  $NH_4Cl$ , 10 mM EDTA, 0.1 mM DFP containing 1 mg of leupeptin and 1 mg of pepstatin. The supernatant obtained by centrifugation was diluted 2-fold with 50 mM Tris HCl, pH 8.0 and 0.1 mM DTT and left for 48 h at 4°C. The sample

was then dialyzed successively against 150 mL, 500 mL, and 2 L of 0.1 M sodium-phosphate, pH 6.0. The dialyzed sample was applied to a DEAE-Sepharose column (1.4 x 18 cm) that was equilibrated with 0.1 M phosphate, pH 6.0. After the column was washed, proteins were eluted by a linear gradient formed by 100 mL each of 0 M and 0.5 M NaCl in the phosphate buffer. The major protein peak that contained anticoagulant activity was pooled and ammonium sulfate was added to 80% saturation. The precipitates collected by centrifugation were dissolved in 5 mL of 50 mM sodium-acetate, pH 5.6, 150 mM NaCl, and 10 mM DTT and applied to a Sephadex G-75 column (2.5 x 90 cm) equilibrated with the same buffer except for DTT. Annexin V-N6 eluted as a single peak and it was homogeneous on SDS-PAGE with a molecular mass of 36 kDa. The yield was 35 mg of protein from 3 L of culture medium. The modified annexin prepared as described above is then used according to the invention.

#### EXAMPLE XI

##### **Preparation of a Modified Annexin V and its Dimer**

Modified annexin V, by addition of accessible sulfhydryl groups, and its dimer, is produced as follows: Transfect the existing plasmid pJ110 (also known as pANXVC-S-N6) into E. coli strain BL21(DE3). Express cytoplasmically by overnight growth to saturation in 4 x 1 L of T broth. Harvest the cells by centrifugation. Sonicate in a buffer consisting of 50 mM TrisHCl pH 7.2,

10 mM  $\text{CaCl}_2$ . Centrifuge (20 min at 23,000 x g), discard supernatant. Release modified annexin from the pellet by suspending in 50 mM TrisHCl pH 7.2, 20 mM EDTA.

Centrifuge (20 min, 23,000 x g), discard the pellet.

5 Treat the supernatant with RNase A (3  $\mu\text{g}/\text{ml}$ ) for 2 h at room temperature. Dialyze in 25,000 molecular weight cutoff tubing against 3 X 2 L of 20 mM TrisHCl pH 8.0 (well aerated); a portion of the modified annexin present is converted to A5 dimer by spontaneous disulfide bond formation due to air oxidation. Purify by ion-exchange FPLC on Pharmacia MonoQ HR10/10 column with a linear gradient from 0 - 0.5 M NaCl in 20 mM TrisHCl pH 8.0. Add 1/30 volume of 1 M (2-[N-morpholino]ethanesulfonic acid; Sigma Chemical) to fractions to maintain pH at 5.8 and inhibit spontaneous disulfide bond formation. The monomer elutes at 0.22 M NaCl and the dimer at 0.28 M NaCl. This procedure yields approximately 5 mg each of pure monomer and dimer per L of starting culture material.

20 Once the disulfide link dimer of modified Annexin V is produced, it can be evaluated for membrane and thrombus binding activity as well as pharmacokinetics and organ uptake in mice. Its affinity for cell membranes is determined by direct titration (1, below), its blood disappearance curve and biodistribution are determined in mice (2, below), and its thrombus:blood ratio determined in vivo in the porcine atrial thrombus model 30 min and 120 min after injection and compared with the

corresponding ratio for monomeric annexin V (3, below).

1. *Binding of modified Annexin V and Annexin V dimer to Phosphatidylserine containing membranes.* If derivatives are not already labeled with Tc-99m, each derivative is labeled with I-125 by the Iodo-Gen method and purified by gel filtration, as previously described for annexin V (see Tait et al, J. Lab. Clin. Med. 123: 741-8, 1994).

The affinity of each derivative for cell membranes is measured by direct titration of preserved erythrocytes with exposed phosphatidylserine as previously described in Tait et al, J. Lab. Clin. Med. 123: 741-8, 1994.

Bound and free ligand are separated by centrifugation of cells through a silicone-oil barrier. The dissociation constant is determined by fitting the data to a simple model of binding to homogeneous sites as described by Tait. Use of artificially treated erythrocytes with high levels of exposed PS is an experimental convenience that gives results equivalent to those obtained with activated platelets with exposed PS. Normal erythrocytes in vivo do not have extracellularly exposed PS (Tait et al, J. Lab. Clin. Med., 123: 741-8, 1994).

2. *Blood clearance, biodistribution, and excretion in mice.* An intravenous bolus of 10  $\mu\text{g/kg}$  (0.3  $\mu\text{g}$  per 30-gram mouse) of each I-125-labeled or Tc-99m-labeled derivative (dimer; galactose-modified annexin; Tc-99m-modified annexin; Tc-99m-seryl succinate-labeled annexin V) is injected. Blood specimens are prepared at 1, 2, 3, 5, 7, 10, 15, 20, 30, 45, 60, 90, and 120 min. The

following organs are harvested from four mice each at 30, 60, and 120 min after injection: brain, heart, aorta, vena cava, lung, spleen, liver, stomach, intestines, kidney, bladder, skeletal muscle, bone, and blood.

Results are expressed as %ID/g tissue and %ID/organ.

3. *Thrombus/blood ratio in vivo in pigs.* Established procedures are used as described for acute atrial thrombi (Stratton et al, Circulation, 92: 3113-21, 1995) for I-125-labeled dimer, galactose-modified annexin, and annexin V as control. Based on previous experience with native annexin V in this model, a twofold difference in thrombus/blood ratios between modified and native annexin V can be demonstrated at a P value of  $\leq 0.05$  in a two-tailed t-test with an n of about 5 animals.

Once a molecule with desirable properties is identified, the annexin V dimer can be produced directly as a fusion protein using an existing expression system as described in Tait et al., J. Biol. Chem. 270: 21594-21599, with the two annexin V moieties connected by a peptide linker. This dimeric molecule can also contain additional functional sites, such as an endogenous Tc-99m chelation site or a free sulfhydryl to allow attachment of galactose.

#### EXAMPLE XII

##### **Endogenous Radiolabeling of Modified Annexin**

A study of Tc-99m radiolabeling of modified annexin (Annexin V-N-6 of Example X) was carried out following the method of George et al, Proc. Natl. Acad. Sci. USA,

92:8358-62, 1995. The modified annexin, 160  $\mu$ g, was combined with 2.5 mCi of Tc-99m pertechnetate, with methylene diphosphonate as the exchange ligand and stannous ion as reductant. The final concentration of the protein as available was 200  $\mu$ g/ml. However, incubation at pH 10 for 30 min at 37°C resulted in 52% radiochemical yield. This apparent yield was consistent with results obtained after purification by gel filtration, which showed recovery of 56% of the applied radioactivity in 89% radiochemical purity. Storage of the purified material indicated no change in radiochemical purity as measured by TLC and Zorbax HPLC chromatography after 18 h in PBS at room temperature. In contrast, native annexin V showed only 2% labeling under the same conditions. Thus, the results have shown that the modified annexin V molecule can be labeled directly with Tc-99m. One of ordinary skill in the art can evaluate the modified annexin V molecule for optimization of Tc-99m radiolabeling yield (protein concentration, exchange chelates, pH dependence, temperature, and time); radiolabel stability (challenge with serum and various concentrations of cysteine); cell and thrombus binding; blood clearance and biodistribution properties; catabolite forms of excreted radioactivity (HPLC of biological samples and comparison with synthetic peptide Tc-99m chelate standards for identification). Suitable methods include those used for characterization of various N,S amide thiolate chelating agent for Tc-99m

bifunctional protein radiolabeling agents (Kasina et al, J. Nucl. Med., 32: 1445-51, 1991; Fritzberg et al, Proc. Natl. Acad. Sci. USA, 95: 4025-9, 1988).


5

### EXAMPLE XIII

#### Production of a Radiolabeled Annexin-Esterase Sensitive Chelate Conjugate



+




10

Annexin-V

pH = 9-10

Time = 15-30 min

25°C-37°C



*Experimental procedure--* Annexin V is radiolabeled with Tc-99m N<sub>3</sub>S-serylsuccinate via the preformed chelate approach as described for antibody Fab fragments (Srinivasan et al, J. Nucl. Med., 32: 1017 (abstract), 1991). Application of this approach with the C<sub>5</sub>N<sub>2</sub>S<sub>2</sub> active ester has been similar for both antibody proteins (Kasina et al, J. Nucl. Med., 32: 1445-51, 1991) and annexin V (Stratton et al, Circulation, 92: 3113-21, 1995). In the present procedure, serylsuccinate ester is substituted in the radiolabeling procedure. This derivative is evaluated in vitro for label stability, membrane binding, and clot binding. It can be injected into mice and its uptake measured in organs including liver and kidney over the period from 10 min to 2 h. This indicates the rate of organ uptake as well as retention, and allows comparison of the time course with previous data for the amide-linked Tc-99m chelate.

Success is indicated by significant reduction in liver retention of radioactivity. Increased rate of organ disappearance following uptake of Tc-99m annexin V indicates the potential of improved thrombus to organ ratios within time of imaging follow up. Imaging experiments can be performed with the porcine model in order to predict comparable improvements under clinical conditions.

*Labeling of annexin V with Tc-99m-serylsuccinate ester.* Annexin V conjugated to the serylsuccinate ester is labeled with Tc-99m according to the OncoTrac kit

procedure (Kasina et al, J. Nucl. Med., 32: 1445-51, 1991), and then conjugated to annexin V using the previously described method for conjugation of Tc-99m-N<sub>2</sub>S<sub>2</sub>-TFP to annexin V (Stratton et al, Circulation, 92: 3113-21, 1995).

#### EXAMPLE XIV

##### **Production of a Radiolabeled Modified Annexin V-Galactose Cluster-Chelate Conjugates**






A maleimide-linked galactose cluster of the present invention can be conjugated specifically to the sulfhydryl of the modified annexin V. After verification of stoichiometry of galactose labeling, the galactose-modified annexin is radioiodinated and evaluated for membrane and thrombus binding in vitro, and pharmacokinetics and organ uptake in mice. Maintenance of binding and faster blood disappearance are followed up by determining in vivo thrombus to blood ratios in the porcine model at 30 and 120 min after injection and compared to corresponding ratios for unmodified annexin V.



The rate of blood disappearance of the modified Annexin V without compromising the ability of the modified annexin V to bind to cell membranes or thrombi in vitro is compared to unmodified annexin V. The in vivo thrombus/blood ratio 30 min after injection should be higher compared to native annexin V, and the same or





higher at 120 min after injection. However, if use of this mechanism also increases the background signal from the liver at the particular times chosen for imaging, use of the esterase-sensitive linker mechanism to accelerate excretion of radiolabel from the liver can be used to decrease the background signal. The following examples represent further embodiments of the present invention.

EXAMPLE XV**Procedure for N-acetylgalactosamine-S-Annexin-N-Chelating  
Compound**


5









5



117




20



22

SMCC/Et<sub>3</sub>N  
DMF, RT




23


1) AG-1 x 8 Anion  
exchange resin  
MeOH/H<sub>2</sub>O, Rt  
17 hrs, filter,  
concentrate





**Ref:**





26

### Preparation of Methyl 6-bromohexanoate (1)

To a 2 liter round bottom flask, charge with 99.7 g (0.511 mol) of 6-bromohexanoate (Aldrich Chemical Co., Milwaukee, Wisconsin) and 1 liter of methanol, was bubbled hydrogen chloride gas for 1-2 minutes. The mixture was stirred at 20-30°C for 18 h and then concentrated via rotary evaporation. The residue was diluted with 500 mL of diethyl ether and washed with 150 mL of de-ionized water, 200 mL of saturated sodium bicarbonate, and then once again with 200 mL of de-ionized water. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated via rotary evaporation. The residue was distilled under vacuum to afford 99.6 g of the product (1) as a colorless oil (93%): b.p. = 93-96°C at 3 mm Hg: <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 3.57 (3H, s), 3.51 (2H, t), 2.30 (2H, t), 1.78 (2H, pentet) and 1.62-1.28 (4H, m) ppm.

Preparation of Methyl 6-Aminohexanoate Hydrochloride (2)

To a 2 liter round bottom flask, charged with 101.3 g (0.722 mol) of 6-aminohexanoate (Aldrich Chemical Co.) in 1 liter of methanol was bubbled hydrogen chloride gas for 3-4 minutes. The mixture was stirred at 20-30°C for 16 h and then concentrated via rotary evaporation. The residue was twice diluted with 500 mL of methanol and re-concentrated (<0.5 mm Hg) to afford 140.1 g of the product (2) as a white solid (100%): <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 9.40 (1H, broad triplet), 3.57 (3H, s), 3.15 (2H, quartet), 2.29 (2H, t), 1.60-1.38 (4H, m) and 1.32-1.19 (2H, m) ppm.

Preparation of Methyl 6-(Trifluoroacetamido)hexanoate (3)

To a 2 liter round bottom flask, charged with 100.2 g (0.552 mol) of amine hydrochloride 2 and 1 liter of methanol was added 100 g (0.703 mol) of ethyl trifluoroacetate followed by 120 mL (0.861 mol) of triethylamine. The mixture was stirred at 20-30°C for 19 h and then concentrated via rotary evaporation. The residue was diluted with 500 mL of diethyl ether and then filtered. The filtrate was washed with 3 x 300 mL aliquots of 1N aqueous HCl, 200 mL of de-ionized water, 2 x 200 mL aliquots of saturated aqueous sodium bicarbonate and finally with 200 mL of de-ionized water. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was distilled under vacuum to afford 115.8 g of the product (3) as a colorless oil: b.p. = 113-116°C at 120 mm Hg: <sup>1</sup>H NMR (d<sub>6</sub>-

DMSO) d 3.57 (3H, s), 2.75 (2H, m), 2.29 (2H, t), 1.60-1.40 (4H, m) and 1.37-1.19 (2H, m) ppm.

Preparation of N,N-Bis-(5-Methoxycarbonylpentyl)amine Hydrochloride (4)

5 To a 5 liter three neck flask equipped with a reflux condenser connected to a gas bubbler, charged with 20.9 g of 60% sodium hydride (0.523 mol) in 1 liter of anhydrous dioxane, was added 100 g (0.416 mol) of secondary amide 3 in 200 mL of dry dioxane over a 20 minute period. The mixture was stirred at 20-30°C for 1 h, and then 130 g (0.622 mol) of bromide 1 in 100 mL of dioxane was added. The mixture was heated to reflux and stirred for 7 h. An additional 10 g of 1 was added and the resulting mixture stirred for 15 h more. The mixture was cooled and concentrated via rotary evaporation. The residue was diluted with 600 mL of 1 N aqueous HCl and extracted with 1 liter of ethyl acetate. The organic phase was then washed with 250 mL of de-ionized water, 250 mL of 5% aqueous sodium metabisulfite, and finally with 250 mL of de-ionized water. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated.

The residue was diluted with 300 mL of de-ionized water and 500 mL of methanol and treated with 200 mL of 10 N aqueous sodium hydroxide. The mixture was stirred at 20-30°C for 16 h and concentrated to a thick syrup via rotary evaporation. The residue was diluted with 800 mL of deionized water and acidified to pH 1-2 with 200 mL of concentrated HCl. The mixture was washed with 3 x 300 mL

5 aliquots of diethyl ether and the aqueous phase then concentrated to a thick syrup via rotary evaporation. The residue was diluted with 1 liter of dry methanol and re-concentrated via rotary evaporation. The residue was diluted with 1 liter of dry methanol and then hydrogen chloride gas was bubbled into the mixture for 2-3 minutes. The mixture was stirred at 20-30°C for 18 h, and then vacuum filtered through Celite (manufactured by J.T. Baker). The solids were rinsed with 200 mL of methanol. The combined filtrates were concentrated. The residue was diluted with 1 liter of methanol and hydrogen chloride gas again bubbled into the mixture for 2-3 minutes. The mixture was stirred for 3 h and then concentrated. The residue was diluted with 1 liter of methanol and 10 g of activated charcoal was added. The mixture was stirred for 30 minutes and then vacuum filtered through Celite. The solids were washed with 100 mL of methanol and the combined filtrates concentrated. The residue was dissolved in hot 2-propanol and then allowed to recrystallize, first at room temperature and then with the use of an ice bath. The solids were filtered and rinsed with 3 x 75 mL aliquots of cold 2-propanol. The solids were air dried to afford 70.5 g of the product (4) as a white solid. The filtrates were combined and concentrated. The residue was recrystallized from 200 mL of 2-propanol to afford an additional 15.3 g of product for a total of 85.8 g (67%): <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 8.69 (2H, broad), 3.57 (6H, s), 2.82

(4H, m), 2.30 (4H, t), 1.67-1.43 (8H, m) and 1.28-1.19 (4H, m) ppm;  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  3.66 (6H, s), 3.42 (4H, t), 2.34 (4H, t), 1.75-1.55 (8H, m) and 1.45-1.25 (4H, m) ppm.

5 Preparation of N-BOC-5-Aminopentanol (5)

To a 2 liter three neck round bottom flask, fitted in the center neck with a 500 mL addition funnel and in a side neck with an adaptor venting to a gas bubbler, was added 40 g (0.388 mol) of 5 aminopentanol in 500 mL of dry acetonitrile. Then 84.5 g (0.387 mol) of di-t-butyl-dicarbonate in 400 mL of dry acetonitrile was added over a 50 minute period. The mixture was stirred at 20-30°C for 15 h and then concentrated. The residue was diluted with 600 mL of ethyl acetate and washed with 2 x 200 mL aliquots of 0.5 N aqueous HCL and 2 x 200 mL aliquots of de-ionized water. The organic phase was dried over anhydrous magnesium sulfate, vacuum filtered, and concentrated, first via rotary evaporation and then using full vacuum pump pressure (<0.5 mm Hg), to afford 74.5 g of the product (5) as a near colorless oil (88%):  $^1\text{H}$  NMR ( $d_6$ -DMSO)  $\delta$  6.72 (1H, broad triplet), 4.31 (1H, t), 3.43-3.27 (2H, m), 2.87 (2H, quartet), and 1.45-1.10 (15H, s and multiplet) ppm;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  4.58 (1H, broad s), 3.65 (2H, t), 3.13 (2H, quartet), and 1.70-1.30 (15H, singlet and multiplet) ppm: Thin Layer Chromatography (Visualization with ninhydrin spray and heat); Silica Gel,  $R_f$  = 0.28 (95/5 methylene chloride/methanol).

Preparation of N-BOC-5-Aminopentyltoluenesulfonate (6)

To a 1 liter round bottom flask, charged with 74.5 g (0.366 mol) of N-BOC-aminopentanol (6) in 400 mL of methylene chloride, was added 45 mL of anhydrous pyridine followed by 74.1 g (0.389 mol) of p-toluenesulfonyl chloride. The mixture was stirred at room temperature for 17 h, diluted with 200 mL of methylene chloride and washed with 400 mL of 0.5 N HCl, 2 x 200 mL aliquots of 0.5 N HCl, and 2 x 100 mL aliquots of de-ionized water. The organic phase was dried over anhydrous magnesium sulfate, vacuum filtered, and concentrated. The residue was chromatographed on 11 x 23 cm of silica gel, eluting first with methylene chloride and then with 3:97 ethyl acetate/methylene chloride. The fractions containing product were combined and concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg), to afford 82.13 g of the product (6) as a white solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.77 (2H, d), 7.31 (2H, d), 4.45 (1H, broad s), 3.98 (2H, t), 3.03 (2H, t), 2.41 (3H, s), and 1.80-1.20 (15H, singlet and multiplet) ppm: Thin Layer Chromatography (Visualization with ninhydrin spray and heat); Silica Gel,  $R_f$  = 0.50 (3:97 ethyl acetate/methylene chloride).

Preparation of 1-b, 3,4,6-Tetra-O-Acetyl-N-Acetyl-Galactosamine (7)

To a 500 mL round bottom flask charged with 25.0 g (116 mmol) of galactosamine hydrochloride (Sigma Chemical Co., St. Louis, Missouri) was added 180 mL of anhydrous pyridine and then 115 mL of acetic anhydride (1.22 mol).

The mixture was stirred at 20-30°C for 44 h and then poured into a 2 liter beaker containing 600 g of ice and 600 mL of de-ionized water. The mixture was stirred at room temperature for 10-15 minutes and then vacuum filtered. The collected solids were rinsed with 4 x 100 mL aliquots of de-ionized water, air dried for 2 h and then dried under full vacuum pump pressure (<0.5 mm Hg) for 14 h to give 39.8 g of the product as a white solid (88%): <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 7.89 (1H, d), 5.63 (1H, d), 5.16 (1H, d), 5.07 (1H, dd), 4.28-3.92 (4H, m), 2.11 (3H, s), 2.02 (3H, s), 1.99 (3H, s), 1.90 (3H, s) and 1.88 (3H, s) ppm.

Preparation of 3,4,6-Tri-O-Acetyl-N-Acetyl-Galactosamine-1-β-Pseudothioure a Hydrochloride (2)

To a 1 liter round bottom flask, charged with 39.8 g (102 mmol) of 7, was added 400 mL of acetyl chloride. The mixture was stirred at 47-48°C for 64 h. The mixture was concentrated and then twice diluted with 200 mL of methylene chloride and re-concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg), to afford 40.2 g of the crude product (8) as a dark amber foamy solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.24 (1H, d), 5.61 (1H, d), 5.43 (1H, dd), 5.27 (1H, dd), 4.83-4.71 (1H, m), 4.48 (1H, t), 4.22-4.01 (2H, 2 dd's), 2.15 (3H, s), 2.02 (3H, s), 2.00 (3H, s) and 1.98 (3H, s) ppm. To the crude chloride (8), in a 1 liter round bottom flask, was added 9.3 g (122 mmol) of thiourea and 150 mL of acetone. The mixture was stirred at reflux for 40

minutes and then cooled in an ice bath for 30 minutes and then vacuum filtered. The collected solids were rinsed with 2 x 75 mL aliquots of acetone. The solids were then air dried for 45 minutes and then dried further under full vacuum pump pressure (<0.5 mm Hg) for 2 h to afford 33.0 g of the product (2) as a light beige solid (74% overall yield from 7):

$^1\text{H}$  NMR ( $d_6$ -DMSO)  $\delta$  9.38 and 9.12 (2 broad s's, 3H), 8.36 (1H, d), 5.56 (1H, d), 5.34 (1H, d), 5.01 (1H, dd), 4.38 (1H, t), 4.22-4.00 (3H, m), 2.11 (3H, s), 2.01 (3H, s), 1.92 (3H, s) and 1.81 (3H, s) ppm.

Preparation of 1-b-Mercapto 3,4,6-Tri-O-Acetyl-N-Acetyl-Galactosamine (10)

To a 1 liter round bottom flask, charged with 30.0 g (67.9 mmol) of the pseudothiurea (2) in 175 mL of methylene chloride and 175 mL of de-ionized water was added 7.08 g (37.24 mmol) of sodium metabisulfite followed by careful addition of 10.2 g (74.5 mmol) of potassium carbonate. The mixture was stirred at room temperature for 40 minutes and the mixture then transferred to a 500 mL separatory funnel. The layers were separated and the aqueous phase was then extracted with 2 x 125 mL aliquots of methylene chloride. The organic extracts were combined, dried over anhydrous sodium sulfate, filtered and concentrated to give 24.2 g (10) of the product as a very pale yellow (off-white) solid (98%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  6.24 (1H, d), 5.61 (1H, d), 5.43 (1H, dd), 5.27 (1H, dd), 4.83-4.71 (1H, m), 4.48

(1H, t), 4.22-4.01 (2H, 2 dd's), 2.15 (3H, s), 2.02 (3H, s), 2.00 (3H, s) and 1.98 (3H, s) ppm.

Preparation of 3,4,6-Tri-O-Acetyl-N-Acetyl-Galactosamine-1-a-S-[5'-Thiopentyl-N-BOC-Amine] (11)

5 To a 1 liter round bottom flask, charged with 24.2 g (66.6 mmol) of the thiol (10) under nitrogen atmosphere, was added 350 mL of dry acetonitrile. The mixture was heated to 40-42°C, the solids eventually dissolving over a 20 minute period. 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU commercially available from Aldrich Chemical Company, 10.5 mL, 70.2 mmol) was then added and the mixture stirred for 20 minutes. Then, 24.0 g (67.1 mmol) of the tosylate 6 in 75 mL of acetonitrile was added over a 3-4 minute period. The resultant mixture was stirred at 40-25°C for 1.5 h and then concentrated. The residue was diluted with 400 mL of methylene chloride and washed first with 250 mL of 0.5 N aqueous HCl and then with 250 mL of 5% aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium, vacuum filtered, and concentrated via rotary evaporation. The residue was chromatographed on 21 x 7 cm of silica gel (manufactured by E.M. Merck), eluting with 55/42.5/2.5 ethyl acetate/hexane/ethanol. The fractions containing product were combined, concentrated and re-chromatographed on 21 x 7 cm of RP-18 silica gel (manufactured by J.T. Baker), eluting with 500 mL each of 50/50, 55/45, 60/40, 65/35, and 70/30 methanol/water and then with 75/25 methanol/water until all of the desired product had

eluted from the column. The fractions containing product were combined and concentrated. The residue was diluted with 500 mL of methylene chloride and treated with anhydrous magnesium sulfate. The mixture was vacuum filtered and the filtrate was concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg) to afford 17.9 g of the product (11) as a foamy white solid: <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 9.38 and 9.12 (2 broad s's, 3H), 8.36 (1H, d), 5.56 (1H, d), 5.34 (1H, d), 5.01 (1H, dd), 4.38 (1H, t), 4.22-4.00 (3H, m), 2.11 (3H, s), 2.01 (3H, s), 1.92 (3H, s) and 1.81 (3H, s) ppm: Thin Layer Chromatography (Visualization with p-anisaldehyde spray and heat); Silica Gel, R<sub>f</sub> = 0.50 (57/40.5/2.5 ethyl acetate/hexane/ethanol); RP-18 Silica Gel, R<sub>f</sub> = 0.21 (65/35 methanol/water).

Preparation of Methyl 6-Methylaminohexanoate Hydrochloride (12)

To a 2 liter three neck round bottom flask, charged with 8.77 g of 60% NaH in mineral oil (219 mmol, 1.1 equiv.) in 500 mL of anhydrous tetrahydrofuran, was fitted a 500 mL addition funnel in the center neck. Then, 34.5 g (144 mmol) of secondary amide 3 in 300 mL of anhydrous tetrahydrofuran was added over a 30 minute period. The mixture was stirred for 50 additional minutes and then 22.6 mL (363 mmol) of iodomethane was added. The mixture was stirred at room temperature for 23 h and then transferred to a 2 liter round bottom flask and concentrated via rotary evaporation. The residue was

treated with 400 mL of 1 N aqueous HCl and then extracted with 300 mL of ethyl acetate and then with 2 x 200 mL aliquots of ethyl acetate. The organic extracts were combined and first washed with 3 x 125 mL aliquots of 5% aqueous sodium thiosulfate and then with 100 mL of de-ionized water. The organic phase was dried over anhydrous magnesium sulfate, vacuum filtered, and concentrated. The residue was dissolved in 250 mL of methanol and re-concentrated. The residue was diluted with 250 mL of methanol and treated with 50 mL of 10 N aqueous sodium hydroxide followed by 100 mL of de-ionized water. The mixture was stirred at room temperature for 17 h, diluted with an additional 50 mL of de-ionized water and then washed with 3 x 200 mL aliquots of hexane. The aqueous phase was concentrated via rotary evaporation. The residue was diluted with 500 mL of methanol and hydrogen chloride gas was bubbled into the mixture for 2-3 minutes (10 g). The mixture was stirred at room temperature for 3 h and then vacuum filtered and concentrated via rotary evaporation. To the residue was added 500 mL of methanol and then hydrogen chloride gas was again bubbled into the mixture for 2-3 minutes (9.2 g). The mixture was stirred at room temperature for 18 h. The mixture was cooled in an ice bath and then vacuum filtered. The filtrate was concentrated by rotary evaporation. The residue was twice diluted with 250 mL of methanol and re-concentrated. The residue was diluted with 300 mL of 2-propanol and treated with 4 g of

activated charcoal for 30 minutes. The mixture was vacuum filtered through Celite and the solids rinsed with 2 x 75 mL aliquots of 2-propanol. The filtrates were combined and concentrated, first via rotary evaporation and then under full vacuum pump pressure. The residue was diluted with 250 mL of methanol and re-concentrated, first via rotary evaporation and then under full vacuum pump pressure. The residue was diluted with 250 mL of methanol and hydrogen chloride gas was bubbled into the mixture for 1-2 minutes (5.0 g). The mixture was stirred at room temperature for 2 h and then concentrated via rotary evaporation. The residue was twice diluted with 250 mL of methanol and concentrated, first via rotary evaporation and finally under full vacuum pump pressure (<0.5 mm Hg) to afford 23.91 g of the product (12) as a very light yellow foamy solid (85%): <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 8.72 (2H, broad s), 3.58 (3H, s), 2.82 (2H, m), 2.49 (3H, s), 2.32 (2H, t), 1.68-1.45 (4H, m) and 1.39-1.21 (2H, m) ppm; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 3.63 (3H, s), 2.97 (2H, t), 2.34 (2H, t), 1.75-1.56 (4H, m) and 1.49-1.31 (2H, m) ppm.

Preparation of N-BOC-N, N-Bis-(5-Methoxycarbonylpentyl)amine (15)

To a 500 mL round bottom flask, charged with 6.34 g (29.1 mmol) of di-*t*-butyl-dicarbonate and 9.00 g (29.1 mmol) of N,N-bis-(5-methoxycarbonylpentyl)-amine hydrochloride (4), was added 125 mL of anhydrous acetonitrile followed by 7.5 mL of triethylamine. The

5 mixture was stirred at room temperature for 22 h and then concentrated via rotary evaporation. The residue was diluted with 300 mL of ethyl acetate and washed with 2 x 100 mL aliquots of 0.1 N aqueous HCl, 100 mL of de-ionized water and 100 mL of 5% aqueous sodium bicarbonate. The organic phase was dried over magnesium sulfate, vacuum filtered and concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg) to afford 10.5 g of product (15) as a near colorless oil (97%): <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO) δ 3.57 (6H, s), 3.07 (4H, t), 2.28 (4H, t), 1.60-1.10 and 1.37 (21H, m and s) ppm; Thin Layer Chromatography (Visualization with ninhydrin spray and heat); Silica Gel, R<sub>f</sub> = 0.33 (20/80 ethyl acetate/hexane); RP-18 Silica Gel, R<sub>f</sub> = 0.17 (70/30 methanol/water).

Preparation of N-BOC-N,N-Bis-(5-Hydroxycarbonylpentyl)amine (16)

20 To a 500 mL round bottom flask, charged with 10.5 g of bis-methyl ester 15, was added 75 mL of methanol followed by 75 mL of 1 N aqueous sodium hydroxide. The mixture was stirred at room temperature for 16 h and then concentrated via rotary evaporation. The residue was diluted with 75 mL of de-ionized water and the pH of the resultant solution adjusted to 2.0-2.5 by slow addition of approximately 75 mL of 1 N aqueous HCl. Then, 200 mL of ethyl acetate was added and the mixture stirred vigorously for 3 minutes. The mixture was transferred to a separatory funnel and the layers separated. The

aqueous phase was extracted with 2 x 150 mL aliquots of ethyl acetate. The organic extracts were combined, dried over anhydrous magnesium sulfate, and vacuum filtered. The filtrates were concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg), to afford 9.52 g of the product as a viscous, nearly colorless oil (98%): <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 3.07 (4H, t), 2.28 (4H, t), 1.58-1.10 and 1.37 (21H, m and s) ppm; Thin Layer Chromatography (Visualization with ninhydrin spray and heat); RP-18 Silica Gel, R<sub>f</sub> = 0.44 (70/30 methanol/water).

Preparation of N-BOC-N,N-Bis-(N',N'-Bis(5-Methoxycarbonylpentyl)-5-Carbamyl pentyl)Amine (17)

To a 1 liter round bottom flask, charged with 9.52 g (27.6 mmol) of bis-acid 16 in 250 mL of anhydrous dimethylformamide, was added 19.0 g (61.3 mmol) of N,N-bis-(5-methoxycarbonylpentyl)amine hydrochloride (4) followed by 30 mL of triethylamine. While the mixture was stirred, 25.7 g (58.1 mmol) of BOP was added. The resulting mixture was stirred at room temperature for 14 h and then concentrated via rotary evaporation. The residue was diluted with 750 mL of ethyl acetate and washed with 250 mL of 0.2 N aqueous HCl, 100 mL of 0.1 N aqueous HCl, 100 mL of de-ionized water, and 2 x 100 mL aliquots of 5% aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, vacuum filtered, and concentrated via rotary evaporation. The residue was chromatographed on 9 x 21 cm of silica gel,

eluting first with 70% ethyl acetate/hexane and then with 100% ethyl acetate. The fractions containing product (17) were combined and concentrated via rotary evaporation. The residue was chromatographed on 7 x 23 cm of RP-18 silica gel, eluting first with 75:25 methanol/water, then with 80:20 methanol/water, and finally with 85:15 methanol/water. The fractions containing product were combined and concentrated, first via rotary evaporation and then under full vacuum pump pressure. The residue was diluted with 500 mL of diethyl ether and the resulting solution was dried with anhydrous magnesium sulfate. The mixture was vacuum filtered and the filtrate was concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg), to afford 17.80 g of product (17) as a near colorless, viscous, oil (75%):  $^1\text{H}$  NMR ( $d_6$ -DMSO)  $\delta$  3.57 (12H, s), 3.18 and 3.07 (12H, 2 t's), 2.32-2.16 (12H, m), 1.61-1.09 and 1.37 (45H, m and s) ppm; Thin Layer Chromatography (Visualization with ninhydrin spray and heat); Silica Gel,  $R_f$  = 0.50 (ethyl acetate); RP-18 Silica Gel,  $R_f$  = 0.30 (85/15 methanol/water).

Preparation of N-BOC-N,N-Bis-(N',N'-Bis(5-Hydroxycarbonylpentyl)-5-Carbamyl pentyl)Amine (18)

To a 500 mL round bottom flask, charged with 7.88 g (9.20 mmol) of the tetramethyl ester (18) in 75 mL of methanol, was added 70 mL of 1 N aqueous sodium hydroxide. The mixture was stirred at room temperature for 16 h and then concentrated via rotary evaporation to

a thick syrup. The residue was diluted with 50 mL of de-ionized water and, with vigorous stirring, the pH of the solution was adjusted to 2-2.5 by slow addition of approximately 70 mL of 1 N aqueous HCl, the product (18) oiling out (one liquid phase separates from another liquid phase) in the process. The mixture was extracted with 200 mL of 3:1 2-propanol/methylene chloride, and then 3 x 100 mL aliquots of 3:1 2-propanol/methylene chloride. The organic extracts were combined, dried over anhydrous magnesium sulfate, filtered and concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg) to afford 7.70 g of a near colorless, thick syrup, consisting (by NMR integration) of 6.93 g of the desired product (18; 94%) and 0.77 g of 2-propanol: <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 3.18 and 3.07 (12H, 2 t's), 2.37-2.12 (12H, m), 1.60-1.10 and 1.37 (45H, m and s) ppm; Thin Layer Chromatography (Visualization with ninhydrin spray and heat); RP-18 Silica Gel, R<sub>f</sub> = 0.50 (70/30 methanol/water).

Preparation of N-BOC-Tet-Gal-NAC-1-a-S-C5 Branch (20)

To a 250 mL round bottom flask, charged with 4.05 g (7.38 mmol) of 3,4,6-tri-O-acetyl-N-acetyl-galactosamine-1-a-S-[5'-thiopentyl-N-BOC-amine] (11), was added 20 mL of methylene chloride followed by 20 mL of trifluoroacetic acid. The mixture was stirred at room temperature for 15 minutes. The mixture was concentrated via rotary evaporation and the residue was thrice diluted with 75 mL of methylene chloride and re-concentrated to

afford 6.27 g of residue, a mixture of desired product (19) and residual trifluoroacetic acid:  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  5.61 (1H, d), 5.41 (1H, dd), 5.01 (1H, dd), 4.62-4.47 (2H, m), 4.11 (2H, d), 2.91 (2H, t), 2.74-2.48 (2H, m), 2.11 (3H, 2s), 2.00 (3H, s), 1.93 and 1.91 (6H, 2s), and 1.37-1.10 (6H, m) ppm. To a separate 250 mL round bottom flask, charged with 1.33 g of the syrup containing 90% 18 by weight (net 1.20 g, 1.50 mmol), was added 50 mL of anhydrous dimethylformamide. In order to remove residual 2-propanol, the mixture was concentrated, first via rotary evaporation and then under full vacuum pump pressure ( $<0.5$  mm Hg). To the residue was added 20 mL of anhydrous dimethylformamide and 10 mL of dry triethylamine. To the resultant, stirred, solution was added a dimethylformamide solution of the crude 19 (in a total of 30 mL of anhydrous dimethylformamide) and the resultant mixture stirred at room temperature for 2 h. The mixture was then concentrated via rotary evaporation. The residue was then diluted with 250 mL of methylene chloride and washed with 2 x 100 mL aliquots of 1 N aqueous HCl, 100 mL of de-ionized water, and then with 100 mL of saturated aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, vacuum filtered, and concentrated via rotary evaporation. The residue was chromatographed on 5.5 x 19 cm of RP-18 silica gel, eluting with 250 mL each of 65:35 methanol/water, 70:30 methanol/water, 75:25 methanol/water, and then with 800 mL of 80:20

methanol/water. The fractions containing product were combined, and concentrated, first via rotary evaporation and then under full vacuum pump pressure to afford 3.55 g of a foamy white solid (94%). This material was then chromatographed on 5.5 x 20 cm of silica gel, eluting with 80:20 ethyl acetate/methanol. The fractions containing only the desired product (20) were combined and concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg), to afford 2.83 g of the desired product as a pure white foamy solid (75%): <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 5.58 (4H, d), 5.42 (4H, dd), 5.01 (4H, dd), 4.63-4.51 (8H, m), 4.20-4.00 (8H, m), 3.35-3.10 (20H, m), 2.73-2.47 (8H, m), 2.32 (4H, t), 2.25-2.08 (20H, m and s), 2.00 (12H, s), 1.93 and 1.91 (24H, 2s), 1.71-1.20 (69H, m and s) ppm; Thin Layer Chromatography (Visualization with ninhydrin spray and heat); Silica Gel, R<sub>f</sub> = 0.47 (75:25 ethyl acetate/methanol); RP-18 Silica Gel, R<sub>f</sub> = 0.33 (80/20 methanol/water).

Preparation of N,N-Bis-(N',N'-Bis(5-Methoxycarbonylpentyl)-5-carbamyl pentyl)-Amine TFA (21)

To a 250 mL round bottom flask, charged with 1.50 g (1.75 mmol) of N-BOC-N,N-bis-(N',N'-bis(5-methoxycarbonylpentyl)-5-carbamyl pentyl)-amine (17) in 15 mL of methylene chloride, was added 15 mL of trifluoroacetic acid. The mixture was stirred at room temperature for 15 minutes and then concentrated. The residue was diluted with 50 mL of methylene chloride and then concentrated via rotary evaporation. The residue

was then diluted with 50 mL of methanol and re-concentrated via rotary evaporation. The residue was again re-diluted with 50 mL of methylene chloride and re-concentrated, first via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg).

Preparation of TFA Salt of Tet-Gal-NAC-1-a-S-Pentyl-amine Branch (22)

To a 100 mL round bottom flask, charged with 790 mg (0.313 mmol) of the N-BOC-Tet-Gal-NAC-1-a-S-C5 Branch (20), was added 10 mL of methylene chloride followed by 10 mL of trifluoroacetic acid. The mixture was stirred at room temperature for 15 minutes and then concentrated via rotary evaporation. The residue was diluted with 150 mL of methylene chloride and washed with 2 x 100 mL aliquots of saturated aqueous sodium bicarbonate. The organic phase was dried over magnesium sulfate, vacuum filtered and then concentrated, first via rotary evaporation and then under full pump pressure (<0.5 mm Hg), to afford 690 mg of the product (22) as a foamy off-white solid (91%).

Preparation of Tet-Gal-NAC-1-a-S-Pentyl Amine conjugate of modified Annexin V via SMCC derivatization (25)

The tetragalactosyl N-AC 1-a-S-Pentylamine (22) is reacted with SMCC reagent in triethylamine and DMF at room temperature. The maleimidyl derivative 23 is triturated with anion exchange resin of the type AG-1 X8 (BioRad; Hydroxide form; 2.6 m equiv./g) in aqueous methanol. The mixture is stirred at room temperature for




15 hours and then vacuum filtered. The resin is rinsed with 50 ml of deionized water and then with 50 ml of methanol. The filtrates are combined and concentrated via rotary evaporation and then under full vacuum pump pressure (<0.5 mm Hg) to afford the product 24 as a dry solid. Conjugation of modified Annexin V-SH with maleimidyl derivatized clustered galactose is carried out in 5-15% DMSO solvent at PH range of 5-8 in borate buffer. The maleimidyl galactose cluster is offered to the modified Annexin V at a molar ratios of 1:1, 5:1 and 10:1. The monomeric clustered galactose-Annexin V construct is then purified by gel filtration techniques. The product is then assayed for sulfhydryl content to determine the degree of derivatization to afford the conjugate 25 in good yield.

Preparation of  $^{99m}\text{Tc}$ -galactose cluster-Annexin V conjugate, 26

The  $^{99m}\text{Tc}$ - $\text{N}_2\text{S}_2$ -TFP ester is prepared by the Verluma™ kit labeling procedure (Kasina, S., Rao, T.N., Srinivasan, A., et. al. *Development and Biologic Evaluation of a Kit for Performed Chelate Technetium-99m Radiolabeling of an Antibody Fab Fragment Using a Diamide Dimercaptide Chelating Agent*, J Nucl Med, 32: 1445-1451, 1991). The  $^{99m}\text{Tc}$  was incorporated into the  $\text{N}_2\text{S}_2$ -ligand via transchelation from  $^{99m}\text{Tc}$ -gluconate by heating. The  $^{99m}\text{Tc}$ - $\text{N}_2\text{S}_2$ -TFP ester is then conjugated with galactose cluster-Annexin V construct in 0.2M bicarbonate buffer, PH 10.0 for 20-30 minutes at room temperature. The  $^{99m}\text{Tc}$ - $\text{N}_2\text{S}_2$ -


galactose cluster-Annexin V conjugate is purified by gel filtration chromatography to afford 30-40% radiochemical yield with a radiochemical purity of  $\geq 85\%$  product 26.

5

EXAMPLE XVI**Procedure for Annexin-S-Galactose-Chelating Compound**

10040503-072004  
10040503-072004





Modified Annexin V CYS-SH  
pH 5-9, RT




Preparation of modified Annexin V monomer conjugate of  
trifunctional lysine adduct containing galactose cluster-  
chelating compound moiety, 36

The epsilon amino BOC protection of the trifunctional lysine adduct of galactose cluster with the chelating compound 31 is reacted with trifluoroacetic acid in methylene chloride. The reaction mixture is stirred at room temperature. The solvent removed under reduced pressure and dried to yield BOC deprotected epsilon amine in good yield. The crude product without further purification is triturated with AG-1 X8 anion exchange resin in aqueous methanol for 20 hours at room temperature. The mixture is filtered, and the resin is

rinsed with methanol. The filtrates are combined and is concentrated to afford the o-deacetylated N-acetylgalactose cluster-chelating compound ligand of the trifunctional lysine as an epsilon amine in good yield. The epsilon amine functionality is reacted with 1.2 equivalents of SMCC reagent in DMF solvent in the presence of 10 equivalents of triethylamine at room temperature. The solvent is removed from the reaction mixture and the crude product is purified on a Silica gel column chromatography to afford the maleimidyl derivatized trifunctional galactose cluster chelating compound-ligand. SMCC adduct of the trifunctional lysine product 36 in good yield.

EXAMPLE XVII**Procedure for Annexin-S-Galactose-(Peptide)-Chelating  
Compound**

1) BOP reagent  
2) 29a, Et<sub>3</sub>N, DMF




1) TFA/CH<sub>2</sub>Cl<sub>2</sub>  
RT  
2) AG-1 X 8 Anion Exchange Resin  
MeOH/H<sub>2</sub>O

[illegible]


↓ SMCC/TEA  
DMF, RT

145



42

Modified Annexin V cys-SH  
pH 5-9, 25°C-37°C



N,N'-Bis-(2-disulfidyl-4-hydroxycarbonylphenyl 4'-hydroxysuccinimidyl carbonylphenyl)-1,3-propyldiamine, 14

N,N'-Bis-(2-disulfidyl-4-hydroxycarbonylphenyl)-1,3-propyldiamine **13** (Also **5** Figure 3) is dissolved in DMF. To the solution, 0.5 equivalents of N-hydroxysuccinimide is added and stirred magnetically. To the reaction mixture, 0.6 equivalents of 10-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide (CDI) as a hydrochloride salt is added. The pH of the reaction mixture is adjusted to between 5 and 6 with Phosphate buffer, 1.0 M, pH 7.0. The progress of the reaction is monitored by thin layer chromatography on silica gel plates. Solvent from the reaction mixture is removed under vacuum and dried. The

crude product is purified on silica gel column chromatography. The fractions containing the mono NHS ester and mono acid are pooled and the solvent removed under reduced pressure and dried to yield the desired compound 14 in low yield.

5     Preparation of Annexin v monomer -S-Conjugate of  
Galactose Cluster-Ligand-of the trifunctional Lysine, 43

10     The N,N<sup>1</sup>-bis(2-disulfidyl-6-hydroycarbonyl phenyl-  
1,3-propyldiamine mono NHS ester (14) is reacted with 1.2  
equivalents of t-butyl tetra glycine carboxylate as its  
hydrochloride salt, commercially available from Aldrich  
Chemical Company, in DMF solvent using 10 equivalents of  
triethylamine as a base. The solvent from the reaction  
mixture is removed under vacuum and the crude product  
purified on silica gel column chromatography to afford  
15     the peptide adduct 38 in good yield. The tertiary butyl  
ester of the compound 38 is removed in TFA-CH<sub>2</sub>Cl<sub>2</sub>  
mixture. The mixture is stirrred for 1 hour and the  
solvent removed and dried to yield the carboxylic acid of  
the ligand-tetra peptide adduct 39 in good yield.

20     The carboxyl functionality of the tetra glycine ligand  
adduct 39 is conjugated with 1.2 equivalents of 3,5,6-tri  
o-acetyl NH acetyl tetra galactosyl e-HNBOC α-amino  
lysine 29a in DMF solvent and 5 equivalents of  
triethylamine in presence of 1.2 equivalents of BOP  
25     reagent. The solvent from the reaction mixture is  
removed under vacuum and dried. The crude product is  
purified by silica gel chromatography to afford compound

40 of the trifunctional lysine epsilon amine protected as BOC.

The BOC group of compound 40 is removed in TFA-CH<sub>2</sub>Cl<sub>2</sub> mixture. The reaction mixture is stirred at room temperature for 1-2 hours. Solvent from the reaction mixture is removed and dried. The crude product is then triturated with AG-1 X8 anion exchange resin in aqueous methanol at room temperature. The mixture is filtered, and the resin is rinsed with methanol. The filtrates are combined and concentrated to afford o-deacetyl-N-acetyl tetra galactosyl ligand-chelating compound of the trifunctional Lysine as an epsilon amine, 41 in good yield. The amino compound 41 is derivatized with 1.2 equivalents of SMCC reagent in 10 equivalents of triethylamine and DMF solvent. The reaction mixture is stirred at room temperature for 1-2 hours. The solvent is removed under reduced pressure and the crude product is chromatographed on silica gel column to yield maleimidyl tetra galactose tetra glycy l peptidyl ligand-chelating compound of the trifunctional Lysine, 42 in good yield. The maleimidyl derivative 42 is then conjugated with modified Annexin V-SH at pH 5-9 in borate buffer and/or phosphate buffer. The reaction mixture is incubated at 25°C-37°C for 2 hours. The sulfur adduct of the modified Annexin conjugate is then purified by gel filtration to give the conjugate 43 in good yield.

EXAMPLE XVIII



Procedure for Dimer-N-Galactose-Chelating Compound

or

S-Annexin-N-Galactose (Endogenous Chelation, no dimer)




or

S-Annexin-N-Galactose-Chelating Compound




1) BOP reagent


2) 22




R = Me, COOH, F, OCH<sub>3</sub>, etc.  
RT, Et<sub>3</sub>N/DMF





1) TFA/CH<sub>2</sub>Cl<sub>2</sub>, RT  
2) AG-1 x 8  
Anion Exchange Resin  
MeOH/H<sub>2</sub>O




1)  Et<sub>3</sub>N, RT

151



$\text{r-Annexin V}$   
 $\text{pH 6-10}$   
 $25^\circ\text{C}-37^\circ\text{C}$

140620-20361600



Preparation of N ε-(tert-butoxy carbonyl) N<sup>1</sup>-α-trifluoroacetyl L-lysine, 28

To a 250ml round bottom flash, charged with 10.0g (0.041 mole) of N ε-(tert-butoxy carbonyl) L-lysine 27, commercially available from BACHEM Bioscience Inc. (subsidiary of BACHEM Switzerland, 3700 Horizon Drive, Renaissance at Gulph Mills, King of Prussia, PA, 19406, USA), and 100ml of methanol is added 10.0g (0.07 mole) of ethyl trifluoroacetate followed by 20.0ml (0.144 mole) of triethylamine. The mixture is stirred at 20-30° C for 20 hours and then concentrated via rotary evaporation. The residue is diluted with 50ml of diethyl ether and then filtered. The filtrate is then washed with 3X 30ml aliquots of 1N aqueous HCL, 20ml of deionized water, and finally with 20ml of deionized water. The organic phase is dried over anhydrous magnesium sulfate, filtered and concentrated. The residue is distilled under vacuum to

yield product 28 in good yield. N  $\epsilon$ -(tert-butoxy carbonyl) N<sup>1</sup>- $\alpha$ -trifluoroacetyl L-lysine is reacted with tri-o-acetyl tet-Gal-NAC 1-a-S-Pentylamine in the presence of a BOP reagent. The activated carboxylic acid reacts with the clustered galactosamine in DMF and Triethylamine to afford galactose cluster lysine adduct 29 in good yield. The N  $\alpha$ -trifluoroacetyl protection of the lysine is removed to afford Primary  $\alpha$ -amino lysine-galactose cluster 29a in moderate yield.

Conjugation of galactose cluster with chelating compound and annexin components, 34

Tetragalactosyl-chelating compound construct, 29a-34

To a mixture of amine extended tetra-galactosyl amine, 29a 1.2 equivalents of N,N<sup>1</sup>-bis (2 disulfidyl-4-hydroxycarbonylphenyl)-1,3-propyldiamine mono NHS ester in dimethylformamide, 5 equivalents of triethylamine is added. The mixture is stirred at 15-30°C for 2-24 hours and then concentrated. The residue is diluted with deionized water, and the pH is adjusted to 2.5 by the addition of 1 N aqueous hydrochloric acid. The mixture is washed with ethyl acetate. The aqueous phase is concentrated. The residue is chromatographed on reverse Phase C-18 silica gel. The fractions containing the product are combined and concentrated to give the product 30 in good yield. The BOC protected epsilon amino functionality of compound 30 is removed by stirring at room temperature in trifluoroacetic acid-methylene chloride mixture. The resulting primary epsilon amino

compound, 31 is triturated with AG-1 X8 anion exchange resin (hydroxide form) in methanol-H<sub>2</sub>O mixture at room temperature for 15 hours. The mixture is filtered, and the resin is rinsed with methanol. The filtrates are combined and concentrated to afford the o-deacetylated N-acetyl galactose cluster which upon reaction with succinic anhydride in triethylamine and DMF at room temperature afforded the hemisuccinic acid adduct 32 in good yield. The hemisuccinic acid functionality of compound 32 is activated via 1.1 equivalents of N-hydroxy-succinimide reaction in presence of 1.2 equivalents of DCC coupling reagent. The reaction product is concentrated in vacuo to yield a crude product which upon purification on silica gel chromatography afford the NHS ester 33 in moderate yield. The NHS ester of the galactose cluster chelating compound adduct of the trifunctional lysine is reacted with Annexin V at PH 6-10 in the presence of DMSO (0-15%). The reaction is carried out at 25°C-37°C for 1-2 hours. The crude conjugate is purified by gel filtration chromatography to afford galactose cluster-chelating compound-Annexin V lysine conjugate 34 in good yield.

EXAMPLE XIXRadiolabeling Procedures for Examples XVI, XVII and XVIIICharacterization of Galactose Cluster-Chelate-Annexin V Conjugates

5 Protein concentration is determined using  $A_{280}$  of 0.6 for 1 mg/mL solution of annexin V. The number of galactose residues per molecule of annexin V is determined by measuring the total number of reactive amines on Annexin V before and after reaction with galactosyl Cluster-Chelate conjugate using trinitrobenzenesulfonic acid, as described by Habeeb, Analytical Biochemistry, 14: 328-36, 1966.



10 Also the number of galactose residues per molecule of modified annexin V conjugated via sulfhydryl is determined by measuring the reactive sulfhydryl on Annexin V before and after reaction with galactosyl cluster chelate conjugate using 3-carboxy-4-mitrophenyl disulfide; DTNB; Ellman's reagent, as described by Deakin, et. al.; Biochem. J., 89: 296, 1963. The ability of galactose-chelate-annexin V to bind to activated platelets is assessed by determining its ability to inhibit the binding of unmodified, I-125-radiolabeled annexin V to freshly isolated human platelets, following the method of Thiagarajan and Tait, J. Biol. Chem., 265: 17,240-43, 1990.


Radiolabeling procedure for use in Post-Formed Chelate  
Conjugation Method.

Method A: Stannous gluconate kits are prepared containing 5 mg sodium gluconate, 100 micrograms stannous chloride, 1.0 mg (1 mg/mL) of galactose cluster-chelate-annexin V, and 0.1 to 1.0 mg of lactose. The pH is maintained between 5 and 7 using HCl, acetic acid or NaOH. To the stannous gluconate kit is added 1.0 mL sodium pertechnetate (Tc-99m) with a specific activity of about 50 mCi. The vial is incubated at 25-30°C for 5-30 minutes. The percent formation of labeled conjugate, remaining pertechnetate, and hydrolyzed/reduced technetium is determined by ITLC in 12% TCA as developing solvent.

Method B: Stannous tartrate kits are prepared in an evacuator vial under nitrogen to contain 0.5 mL of disodium tartrate (10 mg/mL) and 0.1 mL stannous chloride (1.0 mg/mL in ethanol). The pH of the solution is kept between 5 and 7, preferably 6.0. To this stannous tartrate solution is added 1.0 mL of sodium pertechnetate (50 mCi), and the solution is allowed to stand at room temperature. In an evacuated vial, 200  $\mu$ l of sodium phosphate (0.5 M, pH 8 or 10) and 1.0 mL of galactose cluster-chelating compound-annexin V conjugate (1.0 mg/mL) are added successively. Then Tc-99m-tartrate (50 mCi) is added, and the vial is incubated at 25-37°C for 5-30 minutes. The percent formation of labeled conjugate, remaining pertechnetate, and hydrolyzed/reduced

technetium is determined by ITLC in 12% (w/v) trichloroacetic acid as developing solvent.





### EXAMPLE XX

#### Production of Modified Annexin V Dimer by Recombinant Methods

A dimer of two modified annexin V molecules can be prepared by recombinant-DNA methods following the methods described by Tait et al., J. Biol. Chem., **270**: 21594-21599, 1995, for construction and expression of chimeric molecules containing annexin V. First, PCR is performed on the annexin V cDNA template pPAP-I-1.6 with oligonucleotide primers that introduce an NdeI site at the 5' end and a BamHI site at the 3' end of the annexin V coding sequence (amino acids 1-320 + stop codon), and this PCR product is cloned by standard procedures into the NdeI and BamHI sites of plasmid pET-12a (Novagen Corp., Madison, WI) to create plasmid pET-12a-Ax1. PCR

is then performed again on the annexin V cDNA template pPAP-I-1.6 with oligonucleotide primers that introduce an NdeI site at the 5' end and a sequence encoding the amino acids Gly-Gly-Gly-Gly-Gly-Gly followed by an NdeI site at the 3' end of the annexin V coding sequence (amino acids 1-320). This PCR product is then cloned by standard procedures into the NdeI site of plasmid pET-12a-Ax1 to create plasmid pET-12a-Ax2. Production of the dimeric annexin V molecule by cytoplasmic expression in E. coli from plasmid pET-12a-Ax2 is then performed by standard procedures, for example as shown by Tait et al., J. Biol. Chem., 270: 21594-21599, 1995.

All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually incorporated by reference.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.